



Threats and challenges for the conservation of marine molluscs in the southwestern Atlantic

Cristiane Xerez Barroso¹ · Eliane Pintor de Arruda² · Florencia Arrighetti³ · Andrés Averbuj⁴ · Gregorio Bigatti^{4,5} · Ramiro Braga^{6,7} · Alvar Carranza^{8,9} · Susete Wambier Christo¹⁰ · Mariana Osório Côrtes¹¹ · Paulo Márcio Santos Costa¹² · Mariano Cumplido⁴ · Omar Defeo¹³ · Augusto Luiz Ferreira Jr.¹⁴ · Manuel Haimovici¹⁵ · Tatiana Silva Leite¹⁶ · Fabrizio Marcondes Machado¹⁷ · Rafaela Camargo Maia¹⁸ · Inês Xavier Martins¹⁹ · Helena Matthews-Cascon¹ · Marcel Sabino Miranda^{17,20,21} · Nicolás Ortiz^{6,22} · Vinicius Padula²³ · Maria Gabriela Palomo³ · Flávio Dias Passos¹⁷ · Guido Pastorino³ · Natan Carvalho Pedro¹¹ · Pablo E. Penchaszadeh³ · Fabrizio Scarabino²⁴ · Javier H. Signorelli⁴ · Leonardo Santos de Souza²⁵ · Paula Spotorno²⁶ · Lenita de Freitas Tallarico²⁷ · Diego Urteaga³ · Silvana Van der Molen⁶ · Valeria Teso³

Received: 31 July 2024 / Accepted: 29 April 2025

© The Author(s), under exclusive licence to Springer Nature Switzerland AG 2025

Abstract

The southwestern (SW) Atlantic is bordered by the coastlines of Brazil, Uruguay, and Argentina and is characterized by a great diversity of ecosystems, in which estuarine and marine molluscs are among the most representative invertebrates. The accelerating degradation of the environment due to human activity is resulting in the loss of species, leaving us insufficient time to comprehensively understand their morphology, ecological interactions, and the ecosystem services they provide. The aim of this manuscript is to review the conservation status of estuarine and marine molluscs in the SW Atlantic, identify the main threats and knowledge gaps, and indicate the challenges and prospects for their conservation. Physical degradation of environments, pollution, exotic species, and overexploitation of fishery resources are among the main non-climatic drivers threatening marine molluscs in the SW Atlantic. In addition, climate threats, such as rising temperatures and sea levels, and ocean acidification, create synergistic effects that are still not fully understood. Given this scenario, it is imperative to implement measures to mitigate this situation. These measures include strengthening management and governance to conserve habitats and control pollution and non-native species, with the aim of reducing ecosystem deterioration and establishing the foundations for monitoring environmental changes through long-term interdisciplinary projects. Additionally, there is a need to improve legislation aimed at regulating fishery practices. To achieve the effective conservation of estuarine and marine molluscs in the SW Atlantic, it is also essential to strengthen scientific dissemination projects and expand and support the training of researchers specialized in molluscs.

Keywords Biodiversity · South America · Anthropogenic impacts · Climate change · Environmental management

“La mer est le vaste réservoir de la nature”.
Jules Verne, *Vingt Mille Lieues sous les Mers*.

Introduction

The concepts of the Anthropocene and the Great Acceleration show the clear interlinked nature of human socio-economic activities, the planet’s biophysical subsystems or processes, and the mid-twentieth century changes in the state and functioning of the Earth System (Steffen et al. 2015a). In this scenario, a set of frameworks and actions [e.g., the Convention on Biological Diversity of the United Nations Environment Programme, the 2030 Agenda, and their 17 Sustainable Development Goals, and the Decade of Ocean

Extended author information available on the last page of the article

Science for Sustainable Development (2021–2030)] have been developed to help understand the risks associated with these changes and to provide guidance for Earth System Governance (Rockström et al. 2009; Steffen et al. 2015b; United Nations 2024a, b).

The vast richness of biota and the associated knowledge gaps in marine biomes make their conservation a great challenge. The species extinction curve is increasing exponentially [Intergovernmental Science–Policy Platform on Biodiversity and Ecosystem Services (IPBES) 2019; Obura et al. 2023], causing the scientific community to start thinking outside the box, looking for new methods and solutions to bend the curve and to safeguard ecosystems. Studies have already shown the impacts of anthropogenic stressors acting simultaneously on marine populations (e.g., marine pollution, habitat degradation, overfishing, the introduction of non-native species, and climate-change related stressors) (Poloczanska et al. 2013; Pecl et al. 2017; Cooley et al. 2022; Costello et al. 2022).

Molluscs are crucial components of the ecological balance of marine fauna, performing different functions in trophic niches and influencing the biodiversity of coastal and oceanic ecosystems (Gofas and Zenetos 2003). More than 50,000 species of marine molluscs are found throughout the world's oceans (Bouchet et al. 2023), on the ocean bottom or in the water column, from the intertidal to the deep sea, from the poles to the tropical latitudes, and this number is still considered an underestimate (Appeltans et al. 2012). Bouchet et al. (2016) suggested that ca. 150,000 species of marine molluscs remain to be described. Although marine species face significant threats, there are few and, in some cases, controversial, examples of extinction of marine molluscs (e.g., *Lottia edmitchelli*, *Cerithideopsis fuscata*, and *Littoraria flammea*) (Cowie et al. 2022; del Monte-Luna et al. 2023). Nonetheless, there are reasons to suspect that many marine extinctions have been generally overlooked and that several marine species are undoubtedly on the path to risk and extinction (Cowie et al. 2022), probably before being described. Currently, based on the categories and criteria established by the International Union for Conservation of Nature, 36 species of marine molluscs are classified as follows: Critically Endangered (14 species), Endangered (ten species), and Vulnerable (12 species). The low number of species evaluated is probably due to the scarcity of long-term studies that include population and reproductive data, which are more difficult to obtain for these groups compared to other animal groups.

The southwestern (SW) Atlantic, which is bordered by the coastlines of Brazil, Uruguay and Argentina, is characterized by a great diversity of ecosystems (Fig. 1). In its tropical area, there is a heterogeneous reef system, which includes the Great Amazon Reef System, the Brazilian semi-arid coast, and the Eastern Brazilian Reef System

(Leão et al. 2003; Banha et al. 2022; Carneiro et al. 2022). This tropical area is also characterized by the presence of four oceanic islands/archipelagos (the Rocas Atoll, the Fernando de Noronha Archipelago, the São Pedro and São Paulo Archipelago, and the Trindade and Martin Vaz Archipelago) and a variety of ecosystems, including mangrove forests, seagrass beds, rhodolith beds, sandy beaches, rocky shores, and estuaries (Dominguez 2006; Mohriak 2020). The subtropical region is defined by vast sandy beaches interposed with rocky coastlines, with many bays and inlets (Dominguez 2006). The temperate zone features equally extensive sandy shores and, although it lacks or rarely has rocky cliffs, it has an extensive shelf composed of a mosaic of sediments and outcrops (Ab'Saber 2005; Parker et al. 1999). The cold region is characterized by wide tidal flats and high rocky cliffs among gravel and sand beaches (Coronato et al. 2008; Mateucci 2012). The deep-sea area of the SW Atlantic includes a variety of geomorphological features, such as sedimented plains, slopes, plateaus, canyons, seamounts, guyots and methane seeps, which host a great diversity of ecosystems, including those of deep-sea corals (Violante et al. 2017; De Leo et al. 2020; Burone et al. 2021; Carvalho et al. 2023).

Estuarine and marine molluscs in the SW Atlantic remain under-researched with respect to their ecology and species inventories, including basic population, reproductive, and trophic data, with some outstanding ecosystems in particular being poorly studied (e.g., reefs at the mouth of the Amazon River, rhodolith beds, and deep-sea communities); there is also an alarming lack of long-term studies on these organisms (Oliveira and Absalão, 2009; Horta et al. 2016; Corrêa et al. 2018; Santos et al. 2021; Gaurisas and Bernardino 2023). However, there has been a marked increase in malacological studies in the SW Atlantic in recent decades. Machado et al. (2023) compiled a list of 2525 valid species occurring along the Brazilian coast, which represent 71% of all Brazilian molluscs. Over 500 marine and estuarine mollusc species have been reported from Uruguay (Scarabino 2004; Scarabino et al. 2015) and, for the Argentine coast, Bigatti and Signorelli (2018) reported 862 valid molluscan species (approximately 30% of the marine invertebrate fauna recorded in this area).

Estimating and understanding the mollusc biodiversity of the extensive western Atlantic area require more research. The intrinsically high richness and morphological variation of Mollusca, along with the difficulty of collecting and/or finding some groups in situ, and the low number of specialists and scarce funding for biodiversity studies, bring great challenges for this field of taxonomic, systematic, and ecological research (Norman and Huchberg 2005; Leite and Mather 2008; Todt 2013; Lima et al. 2017; Campagnari and Geiger 2018; O'Brien et al. 2021; Santos and Carbayo 2021). In the last decade, studies have




Fig. 1 The studied countries (Brazil, Uruguay, and Argentina) in the southwestern (SW) Atlantic Ocean, with their respective marine protected areas (MPAs) highlighted in red. Map created in QGIS using

data from the Ministry of Environment and Climate Change (Brazil), the Ministry of Environment (Uruguay) and the Ministry of Defense (Argentina)

been conducted on lesser-known groups, such as Scaphopoda (Souza et al. 2013, 2020), Polyplacophora (Urteaga 2011; Güller et al. 2015; Jardim et al. 2022), Aplousobranchia (Ivanov and Scheltema 2008; Passos et al. 2019, 2021, 2022), and mesophotic and deep-water molluscs (Gleadall et al. 2010; Guerra et al. 2011; Signorelli and Pastorino 2015; Pastorino and Sánchez 2016; Signorelli and Crespo 2017; Siegwald et al. 2020; Barroso et al. 2021; Souza et al. 2021; Pacheco et al. 2022; Sánchez and Pastorino 2020). Advances in molecular methodologies have also sought to shed light on potential species complexes and the description of new species (Leite et al. 2008c, 2021; Claremont et al. 2011; Padula et al. 2014; De Biasi et al. 2016; Sales et al. 2019; Avendaño et al. 2020; Fernandes et al. 2021a; Jesus et al. 2021a; Dornellas et al. 2022; García-Méndez et al. 2022).


The objective of this review is to analyze aspects of estuarine and marine molluscs in shallow and deep habitats of the SW Atlantic (5°N to 58°S), focusing on the main threats, measures and ongoing management, challenges, and prospects for achieving their effective biodiversity conservation.

Threats to conservation

Marine biodiversity is threatened by several hazards, which can be classified as primary non-climate drivers (e.g., habitat loss, sea pollution, economic extractive activities, and non-native species) and climate-induced drivers (e.g., a long-term increase in sea surface temperature, acidification, the El Niño Southern Oscillation, and marine heat waves) (Cooley et al. 2022).

These two main types of threats interact with each other at local or regional scales, resulting in effects that can be synergistic, antagonistic or additive (He and Silliman 2019). Regarding the estuarine and marine mollusc species of the SW Atlantic, studies have indicated a set of local and regional impacts, including physical habitat degradation, marine pollution, fisheries/shell harvesting, non-indigenous species, increase in seawater temperature, and ocean acidification, as the main ones (see Fig. 2 and sections below).

Fig. 2 Main threats to estuarine and marine mollusc species of the SW Atlantic Ocean, and some management measures that could help to safeguard their biodiversity. The number of studies related to threats is indicated by colored thermometers (*red* high number of studies, *orange* reasonable number of studies, *blue* low number of studies), and countries where conservation measures have already been implemented are indicated by a *check mark*



Primary non-climate drivers

Physical habitat degradation

The tropical and subtropical areas of the SW Atlantic are home to four of the ecosystems most heavily affected by human activities: sandy beaches, estuaries, mangroves, and coral reefs. The rapid growth of urbanization, fisheries, and tourism activities in Brazil, Uruguay, and Argentina in recent decades, together with the lack of controlled planning for natural coastal resources, has led to the construction of ports and coastal defenses (e.g., harbors and jetties) (Marcomini and López 2006). These structures, together with vehicular and ship traffic, the cultivation of seaweed, crustaceans and molluscs, and the extraction of sea salt, affect the morphodynamics of sandy beaches and estuaries, directly influencing communities of benthic and coastal molluscs (Amaral et al. 2016; Santos and Maia 2021; Corte et al. 2023), such as the bivalves *Amarilladesma mactroides*, *Brachidontes rodriguezii*, and *Donax hanleyanus* (Bom and Colling 2022; Elias et al. 2022). Estuarine areas have historically been favorable sites for human settlement and, more recently, have undergone intense urbanization, with many estuaries suffering from high levels of urban effluents, resulting in eutrophication and declining environmental quality (Barletta et al. 2019). In turn, the deforestation of mangroves areas, caused by shrimp farming, can influence shell morphology, size, and density of the gastropods *Littoraria angulifera* and *Melampus coffea* (Tanaka and Maia 2006; Maia and Tanaka 2007; Maia and Coutinho 2013; Araújo and Maia 2021).

With regard to reef areas, the biodiversity of the Great Amazon Reef System, of which still little is known (it is estimated that less than 5% of the reef area has been researched so far), is currently threatened by large-scale oil and gas exploration (Francini-Filho et al. 2018; Banha et al. 2022; Instituto Internacional Arayara 2023). So far, only a preliminary list of molluscs from this region has been published, which was obtained from bottom trawling (a current threat to this ecosystem) and cephalopod surveys using samples from red snapper stomachs (Sales et al. 2019; Santos et al. 2021). The physical destruction of areas of coral reef may also affect the population of the octopus *Callistoctopus furvus* in Bahia State (northeastern Brazil), where the highest density of this species occurs (Jesus et al. 2021a, b).

Deep-sea ecosystems are also threatened by anthropogenic stressors, especially mining activities. The economic and political pressures to exploit the seafloor run at a faster pace than scientific investigations (Sigwart et al. 2017; Rabone et al. 2023). Currently, the Rio Grande Rise, a large oceanic plateau located in the Economic Exclusive Zone (EEZ) of Brazil, is considered a target area for deep-sea mining (Praxedes et al. 2019). The diversity of molluscs of this region is still little known, and what is known is mainly based on empty shells (Pimenta et al. 2018; Fernandes et al. 2021b) and preliminary surveys (T.S. Leite, personal communication). In the Argentine Sea, resource exploitation initiatives have intensified in recent years, including planned mining activities (M.G. Palomo, personal communication) and the significant expansion of hydrocarbon exploration and extraction—particularly shale gas and oil—in the North

Argentine Basin, prompted by a series of government resolutions issued in 2018 (Socrate et al. 2024).

Marine pollution

Table 1 summarizes the studies carried out in the SW Atlantic, most of which investigated the bioaccumulation of metals and contamination with persistent organic pollutants (e.g., polychlorinated biphenyls, petroleum hydrocarbons and polyaromatic hydrocarbons, butyltin and organochlorine pesticides). Despite the high number of publications on pollutant accumulation, and especially the occurrence of imposex (masculinization of females exposed to organotins such as tributyltin, which has been consistently associated with zones of high marine traffic), there are still few studies that have assessed the biological effects of environmental pollutants through biomarker detection (e.g., oxidative stress, shell structure and composition and histological alterations) (see Table 1).

Regarding contaminants of emerging concern (e.g., pharmaceuticals, cosmetics, endocrine disruptors, biocides, and plastics), research is still scarce (Table 1). Some studies have already shown the presence of microplastics in edible species and in marine protected areas (MPAs) of the SW Atlantic (Nunes et al. 2023a), and preliminary data indicate that depuration can be an alternative method for the mitigation of microplastics contamination in oysters (Christo et al. 2021). However, the biological effects of ingesting molluscs contaminated with microplastics is still a knowledge gap. Other types of interaction with marine debris have been observed in cephalopods (which use it as a shelter) and bivalves (which use it as a means of transport over a long distance, also known as “rafting”) in the SW Atlantic (see Table 1). There is a need for further studies on these interactions, especially concerning the plastics and contaminants of the debris (Freitas et al. 2023).

Besides responses at the individual level, the effects of coastal pollution, with the increasing runoff of contaminated water, have led to the loss of seaweed biodiversity in estuarine and marine areas in the SW Atlantic. This loss affects trophic webs, causing shifts in benthic populations and assemblages (Schnerer et al. 2013; de Oliveira Soares et al. 2016; Martinez et al. 2022). For instance, reductions in macroalgal cover have been associated with decreased functional diversity in mollusc assemblages, likely due to a loss of structural complexity and habitat heterogeneity (Floyd et al. 2020).

Fisheries/shell harvesting

Molluscs from the SW Atlantic have been subjected to intense exploitation by subsistence, recreational, small-scale and/or industrial fisheries (Table 2). Among cephalopods,

the industrial fishery of shortfin squid, *Illex argentinus*, historically represents between 10 and 30% of the total catches in this area (FAO 2022), while other squid species are caught in different types of fisheries. Regarding octopods, some species are targets of industrial (e.g., *Octopus americanus*, which is the most important one) or recreational/artisanal fisheries (e.g., *Octopus insularis*, *Octopus tehuelchus*, and *Enteroctopus megalocyathus*) (Table 2). In Brazil and Argentina, octopus catches reported in the national fishing statistics do not include catches by artisanal fishing nor the identification of the octopus species. The records correspond only to landings of octopus bycatch, or of octopus pot fishery (in the case of Brazil). Some species were tested in aquaculture trials in Brazil and Argentina, but none achieved commercial scale status (Bastos et al. 2018, 2020; Berrueta et al. 2020; Braga et al. 2021, 2022). As octopuses have never been commercially farmed before, there are no laws regulating this practice or the octopuses’ welfare (Andrade et al. 2023).

Bivalve species are also targets of industrial, small-scale and artisanal fisheries, being important sources of food and income for traditional communities of the SW Atlantic (Table 2). An industrial fishery developed targets for the Patagonian scallop *Zygochlamys patagonica*, which has been exploited in Argentine waters since 1996 (Morsan et al. 2017). In the 1970s and 1980s, the scallop *Euvola ziczac* was also a target of industrial fishing (with drags and bottom trawls) off the Brazilian southern and south coasts, but overfishing led to the collapse of the fishery and a drastic decrease in the abundance of this species, from which it has still not recovered (Pezzuto and Borzone 2004). In Brazil, over-exploitation (sometimes acting together with other anthropogenic stressors, such as habitat modification) of natural beds or aggregations are also observed in respect of stocks harvested by artisanal fisheries, e.g., *Perna perna*, *Mytella* spp. and *Crassostrea* spp. (Henriques et al. 2004; Oliveira et al. 2005; Pereira et al. 2001, 2003; Carranza et al. 2011). Population declines are also observed in the yellow clam *Amarilladesma mactroides*, artisanally harvested in Uruguay, and recreationally in Argentina. In Argentina, a decrease in the population of this species has been reported for the beaches of the northeastern coast (Olivier and Penschazadeh 1968; Dadon et al. 2001). In Uruguay, a marked decline in the populations of *A. mactroides* was the result of overfishing between 1980 and 1985 (this species has been commercially exploited since the 1960s) (Defeo et al. 2021), and it was decimated by climate-driven mass mortalities throughout its distribution range during the 1990s (see also the section “Climate-induced drivers”).

Gastropod catches in the SW Atlantic are associated with bycatch in industrial demersal fish fisheries, directed “boom and bust” fisheries, scuba-diving harvesting, baited traps, and subsistence intertidal hand-gathering by traditional

Table 1 Observed impacts of pollution, a primary non-climate driver, on estuarine and marine molluscs of the southwestern (SW) Atlantic Ocean

Observed impacts	Affected species	References ^d
Bioaccumulation of metals	Bivalves (<i>Anomalocardia flexuosa</i> , <i>Aulacomya atra</i> , <i>Brachidontes rodriguezii</i> , <i>Dalocardia muricata</i> , <i>Iphigenia brasiliensis</i> , <i>Mytilus</i> sp., <i>Perna perna</i> , and <i>Phacoides pectinatus</i>) Gastropods (<i>Adelomelon ancilla</i> , <i>Buccinanops cochlidium</i> , <i>Buccinastrum deforme</i> , <i>Nacella magellanica</i> , <i>Odontocymbiola magellanica</i> , <i>Pachycymbiola brasiliana</i> , <i>Tegula patagonica</i> , and <i>Trophon geversianus</i>)	1–18
Contamination with persistent organic pollutants	Bivalves (<i>Ardeamya petitiiana</i> , <i>Aulacomya atra</i> , <i>Brachidontes rodriguezii</i> , <i>Mytilus</i> sp., <i>Perna perna</i> , and <i>Tagelus</i> sp.) Gastropods (<i>Buccinastrum deforme</i> and <i>Siphonaria lessonii</i>)	18–28
Oxidative stress due to accumulation of pollutants	Bivalves (<i>Aulacomya atra</i> , <i>Brachidontes rodriguezii</i> , <i>Mytilus</i> sp., and <i>Perna perna</i>) Gastropods (<i>Nacella magellanica</i> and <i>Siphonaria lessonii</i>)	4, 7–10, 14, 18, 24
Shell structure and composition alterations due to accumulation of pollutants	Gastropods (<i>Buccinastrum deforme</i> , <i>Bostrycapulus odites</i> , <i>Lotia subrugosa</i> , <i>Odontocymbiola magellanica</i> , <i>Siphonaria lessonii</i> , and <i>Stramonita brasiliensis</i>)	29–35
Histological alterations due to accumulation of pollutants	Bivalve (<i>Brachidontes rodriguezii</i>) Gastropod (<i>Siphonaria lessonii</i>)	33, 36
Imposex ^a	Gastropods (<i>Adelomelon ancilla</i> , <i>Adelomelon beckii</i> , <i>Bostrycapulus odites</i> , <i>Buccinanops cochlidium</i> , <i>Buccinastrum deforme</i> , <i>Buccinanops monilifer</i> , <i>Conasprella jaspidea</i> , <i>Hastula cinerea</i> , <i>Leucozonia nassa</i> , <i>Monoplex parthenopeus</i> , <i>Odontocymbiola magellanica</i> , <i>Olivancillaria deshayesiana</i> , <i>Olivancillaria vesica</i> , <i>Olivella minuta</i> , <i>Pachycymbiola brasiliana</i> , <i>Pachycymbiola ferussacii</i> , <i>Pareuthria fuscata</i> , <i>Phrontis vibex</i> , <i>Prunum martini</i> , <i>Pugilina tupiniquim</i> , <i>Stramonita brasiliensis</i> , <i>Stramonita rustica</i> , <i>Thaisella coronata</i> , <i>Trophon geversianus</i> , <i>Voluta ebraea</i> , and <i>Xymenopsis muriciformis</i>)	2, 37–69
Accumulation of tributyltin (TBT) in individuals and egg capsules	Gastropod (<i>Pachycymbiola brasiliana</i>)	2, 70
Contamination with emerging pollutants (e.g., pharmaceuticals, cosmetics, endocrine disruptors, and biocides)	Bivalve (<i>Perna perna</i>)	18, 71–75
Accumulation of microplastics ^b	Bivalves (<i>Amarilladesma mactroides</i> , <i>Aulacomya atra</i> , <i>Brachidontes rodriguezii</i> , <i>Crassostrea tulipa</i> , <i>Limnoperna fortunei</i> , <i>Magallana gigas</i> , <i>Mytella guyanensis</i> , <i>Mytella strigata</i> , <i>Mytilus</i> sp., <i>Perna perna</i> , and <i>Tivela mactroides</i>) Gastropods (<i>Aplysia brasiliana</i> and <i>Nacella magellanica</i>) Cephalopods (<i>Abralia veranyi</i> and <i>Vampyroteuthis infernalis</i>)	76–90
Use of marine debris as a shelter	Cephalopods (<i>Octopus tehuelchus</i> and <i>Paroctopus cthulu</i> ^c)	91–95
Use of marine debris as a means of transport (rafting)	Bivalves (<i>Isognomon bicolor</i> and <i>Pinctada imbricata</i>) Gastropod (<i>Petalocochus varians</i>)	96–99

^aMasculinization of females exposed to organotins such as tributyltin, a specific kind of contamination with organic pollutants

^bA specific kind of emerging pollutant

^cA new species documented only on marine debris, which also used it to brood its eggs

^d1—Gil et al. 2006; 2—Cledón et al. 2006; 3—Giarratano and Amin 2010; 4—Comoglio et al. 2011; 5—Conti et al. 2011; 6—Conti et al. 2019; 7—Duarte et al. 2011; 8—Giarratano et al. 2011; 9—Giarratano et al. 2014; 10—Di Salvatore et al. 2013; 11—Buzzi et al. 2017; 12—Primost et al. 2017; 13—Bigatti et al. 2018; 14—Ruiz et al. 2018; 15—Barbosa et al. 2019; 16—Cledón et al. 2021; 17—Francioni et al. 2004; 18—Santos et al. 2022; 19—Massara Paletto et al. 2008; 20—Arias et al. 2010; 21—Commendatore et al. 2015; 22—Oliva et al. 2015; 23—Laitano et al. 2016; 24—Laitano and Fernández-Gimenez 2016; 25—Primost et al. 2018; 26—Francioni et al. 2005; 27—Francioni et al. 2007a; 28—Francioni et al. 2007b; 29—Laitano et al. 2013; 30—Márquez et al. 2011; 31—Márquez et al. 2017; 32—Harayashiki et al. 2021; 33—Landro et al. 2021; 34—Gouveia et al. 2022a; 35—Gouveia et al. 2022b; 36—Arrighetti et al. 2019; 37—Castro et al. 2000; 38—Penchaszadeh et al. 2001; 39—Fernandez et al. 2002; 40—Camilo et al. 2004; 41—Castro et al. 2004; 42—Bigatti and Penchaszadeh 2005; 43—Castro et al. 2005a; 44—Castro et al. 2005b; 45—Castro et al. 2007a; 46—Castro et al. 2007b; 47—Castro et al. 2007c; 48—Queiroz et al. 2007; 49—Castro et al. 2008; 50—Meirelles et al. 2008; 51—Bigatti et al. 2009; 52—Cardoso et al. 2009; 53—Teso and Penchaszadeh 2009; 54—Averbuj and Penchaszadeh 2010; 55—Arrighetti and Penchaszadeh 2010; 56—Gomes 2010; 57—Lima-Verde et al. 2010; 58—Azevedo et al. 2012; 59—Castro et al. 2012; 60—Teso et al. 2012; 61—Rossato et al. 2014; 62—Petracco et al. 2015; 63—Primost et al. 2015; 64—Rossato et al. 2016; 65—Averbuj et al. 2018b; 66—Maciel et al. 2018; 67—Rodrigues et al. 2020; 68—Ribeiro-Brasil et al. 2021; 69—Santos and Maia 2023; 70—Goldberg et al. 2004; 71—Fontes et al. 2018; 72—Cortez et al. 2018; 73—Cortez et al. 2019; 74—Ortega et al. 2019; 75—Fonseca et al. 2020; 76—Santana et al. 2016; 77—Fernández Severini et al. 2019; 78—Pazos et al. 2020; 79—Pérez et al. 2020; 80—Ríos et al. 2020; 81—Ojeda et al. 2021; 82—Truchet et al. 2021; 83—Vieira et al. 2021; 84—Christo et al. 2021; 85—Staichak et al. 2021; 86—Costa et al. 2022; 87—Ferreira et al. 2022; 88—Nunes et al. 2023b; 89—Ribeiro et al. 2023; 90—Schuab et al. 2023; 91—Alves and Haimovici 2011; 92—Storero et al. 2013; 93—Leite et al. 2021; 94—Freitas et al. 2022; 95—Freitas et al. 2023; 96—Breves et al. 2014; 97—Marques and Breves 2014; 98—Póvoa et al. 2021; 99—Breves and Skinner 2014

communities (Table 2). While the utilization of molluscs as a food source is relatively common along the Uruguayan and Argentine coasts, it is not common in Brazil (although some Brazilian traditional communities collect them specially for subsistence) (Table 2). The most popular use of gastropods in Brazil is for ornamental purposes, and demand for their shells is intense due to an increase in tourism (Dias et al. 2011; Mota et al. 2020). Large-scale shell collecting for reasons other than food was included among the possible, yet to be proven, causes for the local extinction of *Olivancillaria contortuplicata* in Barra del Chuy and La Coronilla (Uruguay) (Scarabino 2004). Along the SW Atlantic coast, gastropods are consumed without official records on their landings.

Non-indigenous species

In recent decades, reviews and new records of non-indigenous molluscan species in the SW Atlantic have been published (Pastorino et al. 2000; Orensanz et al. 2002; Carranza et al. 2010; Spotorno-Oliveira et al. 2018, 2020; Cavaleiro et al. 2019; Gernet et al. 2019; Messano et al. 2019; Amaral et al. 2020; Darrigran et al. 2020, 2025; Schwindt et al. 2020; Teixeira and Creed 2020; Lima and Passos 2021; Soares et al. 2022; dos Santos et al. 2023; Pedro et al. 2023). According to these studies, the non-indigenous molluscan species detected up to now in the SW Atlantic are the gastropods *Eualetes tulipa*, *Indothais lacera*, and *Nassarius foveolatus* (Brazil), *Pleurobranchaea maculata* (Argentina) and *Rapana venosa* (Brazil, Uruguay, and Argentina), and the bivalves *Leiosolenus aristatus*, *Mytilopsis* cf. *sallei*, *Mytilopsis leucophaeata*, *Perna viridis* and *Saccostrea cucullata* (Brazil), *Isognomon bicolor* (Brazil and Uruguay), *Magallana gigas* and *Talonostrea talonata* (Brazil and Argentina).

So far, few impacts on native molluscs have been reported (Carranza et al. 2023). Studies with *I. bicolor* have already shown high densities of this species on intertidal rocky shores along the Brazilian coast that have caused changes in the food habits of the gastropod *Stramonita* sp. (which previously preyed on *P. perna*) (López et al. 2010), with the potential to modify the composition and abundance of the local macrofauna community (Domaneschi and Martins 2002; Fernandes et al. 2004; Breves-Ramos et al. 2010; Martinez 2012; Dias et al. 2013). Living specimens of *I. bicolor* were also found in floating debris washed up on the coast of Uruguay, providing evidence that this species can be dispersed by rafting (Breves et al. 2014). *Perna viridis*, a native species of the Indo-Pacific Ocean, was recently recorded off the southeastern Brazilian coast (Messano et al. 2019; dos Santos et al. 2023). There is no record of impacts of this species on native communities, contrary to observations documented for the Caribbean and North America (Dias et al. 2018). Regarding non-indigenous oyster species, impacts of

S. cucullata (which is native to the Indo-Pacific) on Brazilian benthic communities have yet to be observed, despite their sympatry with native oyster species along the southeastern and southern coasts of Brazil (Amaral et al. 2020). On the other hand, the introduced Pacific oyster *M. gigas* now forms extensive beds on stretches of rocky shores of the Argentine coast, altering the structure of intertidal ecosystems (Mendez et al. 2015; Horta et al. 2022). It has also been found that *T. talonata* may outcompete *Crassostrea tulipa*, and is thus considered a nuisance species for oyster culture (Cavaleiro et al. 2019). *Leiosolenus aristatus* caused damage to the shells of cultured scallop *Nodipecten nodosus* in a marine farm in Brazil, including severe scarring, other deformations, and even death (Simone and Gonçalves 2006).

Among non-indigenous gastropods, the predatory activities of *R. venosa* and *I. lacera* may be affecting the ecological properties of their habitat, such as mussel coverage on rocky bottoms (Carranza et al. 2010; Spotorno-Oliveira et al. 2020; Pedro et al. 2023). However, no studies have quantified the extent of this presumably significant environmental impact. The presence of high densities of the sea slug *P. maculata* on the Argentine coast, where it appears to have become successfully established, may also endanger native species, adversely impacting the local biodiversity and posing potential health risks for humans (due to the possible presence of neurotoxins), particularly in tourist hot spots such as north Patagonian gulfs (Bökenhans et al. 2019, 2022; Battini et al. 2021).

Two species previously registered as non-indigenous on the Brazilian coast, the bivalve *Perna perna* and the nudibranch *Bulbaeolidia alba*, do not have this status confirmed herein (Teixeira and Creed 2020; Darrigran et al. 2020). The former was classified as cryptogenic due to divergences in the results of studies carried out on the origin of this species (Teixeira and Creed 2020; Calazans et al. 2021). The latter, in turn, is not considered cosmopolitan but a complex of species (Carmona et al. 2017), and the Brazilian specimens are possibly of the species *Bulbaeolidia oasis*, originally described from Cuba.

The introduction of non-indigenous species of other taxa to the SW Atlantic and their influence on the molluscan community must also be considered. For example, the green crab *Carcinus maenas* from Europe, which has established non-indigenous populations in Argentina, is a potential invader in southern Brazil (Young and Elliott 2019). This invasive crab has already been observed preying on native molluscan species including the mussel *Perumytilus purpuratus* (one of the main structuring species that affords shelter), and the gastropods *Buccinastrum deforme* (in one of the most important nearshore marine gastropod fisheries of Patagonia) and *Tegula patagonica* (Cordone et al. 2022; Márquez et al. 2024). The invasion of the lionfish *Pterois volitans* into northeastern Brazil (Maggioni et al. 2023) could also affect

Table 2 Mollusc fisheries in the SW Atlantic Ocean

Class	Species	Local	Capture methods	References ^c
Cephalopoda (squids)	<i>Illex argentinus</i>	From southern Brazil and Uruguay to Islas Malvinas (Argentina)	Target of industrial jigging and trawl fisheries; bycatch of bottom trawlers hake fisheries (in the upper slope and continental shelf); bycatch by bottom trawling (on the outer continental slope); bycatch of gillnet fishing directed towards monkfish	1–7
	<i>Doryteuthis pleii</i>	Southeastern Brazil	Bycatch of shrimp fisheries; target of small-scale fisheries; artisanal hand-jigging fisheries	1, 2, 8, 9
	<i>Doryteuthis sanpaulensis</i>	Southeastern Brazil, Uruguay, northern Patagonian Shelf (Argentina)	Bycatch of coastal shrimp bottom trawling; artisanal hand-jigging fishery; artisanal fishing (bycatch of beach seine)	1, 2, 10–13
	<i>Doryteuthis gahi</i>	Patagonian Shelf and around the Islas Malvinas (Argentina)	Specialized bottom trawl fishery; artisanal fishing (bycatch of beach seine); bycatch of shrimp and prawn bottom trawling	5, 10, 12
	<i>Doryteuthis surinamensis</i> , <i>Eucoleoteuthis luminosa</i> , <i>Histioteuthis corona</i> , <i>Hyaloteuthis pelagica</i> , <i>Lolliguncula brevis</i> , <i>Ommastrephes barrami</i> , <i>Ornithoteuthis antillarum</i> , <i>Pholidoteuthis adami</i> , <i>Pickfordioteuthis pulchella</i> , <i>Septoteuthis sepioidea</i> , <i>Sthenoteuthis pteropus</i> , <i>Thysanoteuthis rhombus</i> , <i>Todarodes filippovae</i> ^a	Brazilian coastal shallow or deep waters	Bycatch of trawling fisheries	1, 3, 14–16
	<i>Martialia hyadesi</i>	Southern Patagonia around Islas Malvinas and Georgias del Sur (Argentina)	Jigging over continental slope	5, 17, 18
	<i>Moroteuthopsis ingens</i> ^a	East of Islas Malvinas and international waters of Argentina	Bycatch of hake and squid trawlers on the continental shelf and upper slope	19, 20

Table 2 (continued)

Class	Species	Local	Capture methods	References ^c	
Cephalopoda (octopus)	<i>Octopus americanus</i>	Southeastern and southern Brazil and Uruguay	Target of industrial pot fishery; bycatch of shrimp trawling; by artisanal fishers using harpoons during freediving (off rocky shores)	21, 22	
	<i>Octopus insularis</i>	northeastern Brazil and Brazilian oceanic islands	Target of a commercial pot fishery; by artisanal fishers through freediving and with a hook walking on intertidal reefs (called “polvejamento”)	23–27	
	<i>Eledone gaucha</i> ^a	Southern Brazil	Bycatch of shrimp trawling and pot fishery	1–3, 21	
	<i>Eledone massyae</i> ^a	Southeastern Brazil, coast of the Buenos Aires province and the northern part of Patagonia Shelf (Argentina)	Bycatch of fish, shrimp and prawn trawling fisheries	1, 28, 29, 30	
	<i>Callistoctopus furvus</i> ^a	northeastern Brazil	Artisanal fishery through <i>polvejamento</i> (octopus fishing) (on shallow reefs)	26, 31	
	<i>Robsonella fontianiana</i> ^a	Patagonian coast	Bycatch of octopus by artisanal fisheries	32	
	<i>Octopus tehuelchus</i>	Patagonian coast, in Río Negro and Chubut (Argentina)	Use of hooks and traps in the intertidal and traps in shallow subtidal areas; bycatch in trawling fisheries; bycatch from trawl fishing for shrimp and fish	1, 33–37	
	<i>Enteroctopus megalocyathus</i>	Patagonian coast, in Río Negro, Chubut and Santa Cruz provinces, and Beagle Channel (Argentina)	Use of hooks in the rocky intertidal and subtidal hard bottom areas (by diving in the subtidal); bycatch in trawling fisheries; bycatch in trap fishing for southern king crab zone	33, 38–41	
	Bivalvia	<i>Amarilladesma mactroides</i>	Uruguay and Argentina	Artisanal fishery	42, 43
		<i>Donax hanleyanus</i>	Argentina	Recreational	43
		<i>Zygochlamys patagonica</i>	Northern Atlantic Patagonian gulfs (Argentina)	Industrial trawling fisheries	44
		<i>Aequipecten tehuelchus</i>	Natural Protected Area Peninsula Valdés and the southern part of San Matías gulfs (Argentina)	Small-scale fisheries	45, 46
		<i>Aulacomya atra</i> , <i>Eucallista purpurata</i> , <i>Mytilus platensis</i> , <i>Panopea abbreviata</i>	Northern Atlantic Patagonian gulfs (Argentina)	Small-scale fisheries; harvested by divers (<i>P. abbreviata</i>)	12, 44, 47–50
<i>Anomalocardia flexuosa</i> <i>Evola ziczac</i> ^b		Northeastern and southeastern Brazil Southeastern and South Brazil (São Paulo, Paraná, and Santa Catarina States)	Artisanal fishery Industrial fishing with drags and bottom trawls	51–57 58	

Table 2 (continued)

Class	Species	Local	Capture methods	References ^c
Gastropoda	<i>Adelomelon ancilla</i> , <i>Coronium acanthodes</i> , <i>Fustirriton magellanicus</i> , <i>Odontocymbiola magellanica</i>	Uruguay, Argentina	Bycatch in the industrial demersal fish fisheries	59–62
	<i>Adelomelon beckii</i> , <i>Pachycymbiola brasiliana</i> , <i>Tonna galea</i> , <i>Zidona dufresnii</i>	Uruguay, Argentina	Directed “boom and bust” fisheries	61, 63–69
	<i>Buccinastrum deforme</i> , <i>Buccinanops cochlidium</i> , <i>Nacella magellanica</i> , <i>Odontocymbiola magellanica</i>	Patagonian Atlantic coast (Argentina)	Scuba-diving harvesting, baited traps, intertidal hand-gathering	12, 61, 62, 70–72
	<i>Strombus pugilis</i>	Southeastern Brazil (Rio de Janeiro State)	Artisanal fishing by traditional communities	73

^aSpecies caught in small numbers or accidentally

^bSpecies for which the fishery collapsed due to overfishing

^c1—Costa and Haimovici 1990; 2—Haimovici and Perez 1991; 3—Perez et al. 2004; 4—Perez et al. 2009; 5—Arkipkin et al. 2015; 6—Brunetti et al. 1999; 7—Crespi-Abril et al. 2008; 8—Perez et al. 2002; 9—Judkins and Vecchione 2013; 10—Barón and Ré 2002; 11—Vidal et al. 2013; 12—Elias et al. 2019; 13—Braga et al. 2019; 14—Haimovici et al. 2007; 15—Boletim Estatístico da Pesca e Aquicultura 2011; 16—Rosa et al. 2013; 17—González and Rodhouse 1998; 18—Agnew et al. 2005; 19—Lapikhovskiy et al. 2010; 20—Rosas-Luis et al. 2014; 21—Ávila-da-Silva et al. 2014; 22—Cortes 2022; 23—Leite et al. 2008a; 24—Leite et al. 2008b; 25—Leite et al. 2009; 26—Haimovici et al. 2014; 27—Sauer et al. 2021; 28—Pérez and Haimovici 1991; 29—Ré and Taylor 1981; 30—Ré 1998b; 31—Jesus et al. 2022; 32—Ortiz and Ré, 2011; 33—Ré 1998a; 34—Narvarte et al. 2006; 35—Storero 2010; 36—Bocco et al. 2019; 37—Sánchez-Carnero et al. 2022; 38—Cinti et al. 2003; 39—Ortiz 2009; 40—Ortiz et al. 2011; 41—Ortiz and Ré 2019; 42—Olivier and Penchaszadeh 1968; 43—Dadon et al. 2001; 44—Morsan et al. 2017; 45—Orensanz et al. 2003; 46—Orensanz et al. 2007; 47—Morsan 2003; 48—Orensanz et al. 2006; 49—Narvarte et al. 2007; 50—Morsán et al. 2010; 51—Nishida et al. 2004; 52—Nishida et al. 2006; 53—Gil et al. 2007; 54—Botelho and Santos 2005; 55—Silva et al. 2007; 56—Pezzuato et al. 2010; 57—Nascimento et al. 2022; 58—Pezzuato and Borzone 2004; 59—Carranza 2006; 60—Carranza and Horta 2008; 61—Bigatti and Ciocco 2008; 62—Bigatti et al. 2015; 63—Arrighetti et al. 2011; 64—Riestra et al. 2000; 65—Giménez et al. 2005; 66—Defeo et al. 2011; 67—Marín et al. 2020; 68—Sánchez et al. 2012; 69—Navarro et al. 2019; 70—Nieto Vilela et al. 2019; 71—Cumplido and Bigatti 2020; 72—Cumplido et al. 2022; 73—Moscatelli 1987

the native molluscs there, including the *Octopus insularis* population, since they co-occur and could compete for prey (Leite et al. 2016; Peake et al. 2018).

Climate-induced drivers

Studies have been carried out on the impacts of climate change (increased sea surface temperature and ocean acidification) in the SW Atlantic, which stands out as one of the largest and most significant hot spots for ocean warming worldwide (Ortega et al. 2016; Franco et al. 2020; Gianelli et al. 2021).

Manipulative laboratory experiments conducted with estuarine littorinids (*Littoraria angulifera* and *Littoraria flava*) from the Brazilian coast showed an increase in mortality, weight loss, and changes in the behavior of snails due to an increase in water temperature (Maia and Troncoso 2022). In recent research on species from southern Brazil, Uruguay, and northern Argentina, the yellow clam *Amarilladesma mactroides* showed the greatest sensitivity to climate change among the molluscs evaluated, while the cephalopods *Illex argentinus* and *Doryteuthis sanpaulensis* were among the least sensitive (Gianelli et al. 2023). In the Nuevo Gulf (Argentina), mass mortalities of scorched mussels (*Brachidontes rodriguezii* and *Perumytilus purpuratus*), which were probably related to an anomaly in the sea surface temperature, where it increased by more than 1 °C, were detected in different sites where their coverage, which was close to 90% before the 2019 austral summer, was replaced by bare soil and algae (Mendez et al. 2021). Studies also showed that an increase of sea surface temperature may change the geographic distribution and ecological niche of cephalopod species of the SW Atlantic. For example, *Argonauta nodosus* and *Doryteuthis pleii* have undergone a distributional shift toward the poles (Vales et al. 2017; Martins and Juanicó 2018). In addition, species distribution models predict that *Octopus americanus* will experience a poleward expansion alongside a range contraction in lower Atlantic latitudes (Borges et al. 2022). Meanwhile, for *O. insularis*, niche space expansion from the tropical Atlantic to the temperate northern Atlantic and temperate South America waters is expected (Lima et al. 2020), and this species was recently reported off São Paulo State (Brazil), which means its distribution has expanded 5° south (Leite et al. 2024). Increasing sea temperatures are also affecting the distribution of species of the genus *Mytilus*, causing more heat-tolerant species to invade the areas of distribution of cold-water species, while the latter are migrating towards the poles (Beyer et al. 2017).

Regarding the effects of ocean acidification, the manipulative laboratory experiments conducted by Maia and Troncoso (2022) also showed a decrease in shell size, resulting from the dissolution of the spire. Although there is no

published information about the direct effects of ocean acidification on octopus species in the SW Atlantic, an indirect effect could be responsible for the reduction in the number of empty shells available as shelter to the smaller octopus species that have been found using marine debris as alternative dens (Iribarne 1990; Leite et al. 2021; Freitas et al. 2022).

A few studies only have documented the synergistic effects of primary non-climate and climate-induced drivers in the SW Atlantic. The collapse of *Nodipecten nodosus* sea farming in southern Brazil has been attributed to the negative effects of water pollution (high levels of nutrients and bacteria), combined with warming seawater (Thompson et al. 2023). Another consequence of these joint effects is the increase of frequency and duration of harmful algal blooms of marine coastal environments in recent decades, as a result of global climate change and anthropogenic eutrophication, which have negatively impacted *Amarilladesma mactroides* fishing activities, reducing fishers' revenues and leading to an increase in economic uncertainty (Gianelli et al. 2021). Mass mortality events of *A. mactroides* coincided with the onset of a climate shift (ocean warming) during the 1990s and early 2000s, occurring sequentially in a poleward direction from southern Brazil to Argentina as tropical waters moved poleward (Ortega et al. 2016 and references therein).

Oyster reefs and beds, which are dominant structures in estuaries around the globe, are another ecosystem threatened by multiple stressors and their synergistic effects (e.g., extensive harvesting of wild populations, dredging and trawling, introductions of non-native oysters, diseases, alterations of shorelines, changes in freshwater inflows, and increased loadings of sediments, nutrients, and toxins) (Beck et al. 2011). In a global analysis, oyster reef conditions in the SW Atlantic were categorized as “good” (less than 50% lost) or “fair” (50–89% lost), based on the percentage of the abundance compared to historical levels of abundance (Beck et al. 2011). Similar to oyster reefs, mussels also play a crucial role as ecosystem engineers, with their decline considered critical in the context of environmental change (Borthagaray and Carranza 2007; Carranza et al. 2009).

Conservation measures and management in progress

Some management measures have been undertaken in the SW Atlantic to safeguard mollusc diversity and maintain ecosystem health and services, such as the updating of species inventories, assessing the conservation status of species, developing environmental education projects, and undertaking measures to regulate fisheries (Fig. 2).

Biodiversity inventories

For the elaboration of public policies, decision-making, conservation plans, and effective management, it is indispensable to know how many valid taxa are represented in a study area and how vulnerable the species that are present are.

In Brazil, a group of more than 50 taxonomists maintains the list of mollusc species that is constantly updated as part of the Taxonomic Catalog of the Brazilian Fauna, a governmental open access database (<http://fauna.jbrj.gov.br>) (Machado et al. 2023). In Uruguay and Argentina, several studies, including recent oceanographic expeditions, have been conducted to improve knowledge on mollusc diversity and distribution in these regions (Pastorino and Chiesa 2014; Signorelli and Pastorino 2015; Zelaya 2015; Scarabino et al. 2015; Pastorino 2016b, 2016a, 2019; Pastorino and Sánchez 2016; Bigatti and Signorelli 2018; Di Luca and Pastorino 2018; Siegwald et al. 2020; Sánchez and Pastorino 2020; Pacheco et al. 2022, 2024; Cetra and Roche 2023; Sánchez et al. 2023, 2024, 2025).

Although the increasing number of studies undertaken in these areas enhances our knowledge about mollusc biodiversity there, the use of empty shells in taxonomic and biodiversity research hampers our understanding of the true diversity of extant species, as these assemblages could have been on the seafloor for centuries or millennia, and may represent rare extant or extinct taxa (Kowalewski et al. 2023). These factors must be taken into consideration in future studies.

Species conservation assessments

In the most recent Brazilian assessment, based on the categories and criteria established by the International Union for Conservation of Nature, four mollusc species were considered under threat (cephalopods were evaluated separately); of these, the three gastropods were classified as Vulnerable (*Petaloconchus myrakeenae*, *Macrostrombus costatus* and *Titanostrombus goliath*) and the bivalve as Endangered (*Euvola ziczac*) (ordinance ICMBio no. 148, 7 June 2022). The vermetid *P. myrakeenae*, categorized as Vulnerable, was recently proposed as a junior synonym for *Petaloconchus varians* (Breves et al. 2022), and therefore its conservation status should be reviewed. In relation to cephalopods, in the first Cephalopods Threatened Species Evaluation List for Brazil, published in 2024 (<https://salve.icmbio.gov.br/#/>), six species (three octopuses and three squids) were classified as Data Deficient and 40 species as Least Concern. The Argentine Malacological Association is set to publish the red book of molluscs from Argentina [*Libro Rojo de Moluscos Argentinos: Categorización de la Malacofauna según su Estado de Conservación*] (in press), which will be the first assessment and compilation of vulnerable molluscs in the country.

Since there are still limited data on the diversity and ecology of many estuarine and marine mollusc species, the use of life history traits, such as body size and trophic position, can be a useful preliminary screening criterion for assessing a species' vulnerability to extinction. A recent study on benthic gastropods in the western Atlantic showed a negative association between large species and their trophic position, suggesting potential energy restriction and heightened vulnerability to environmental changes (Carranza and Arim 2023). These results indicate the importance of employing screening methods (such as the use of life history traits) prior to regional or global conservation assessments for mollusc species.

Environmental education projects

Environmental education can play a role in increasing public awareness about conservation, especially when conducted in collaboration with relevant environmental authorities. These educational initiatives usually have a localized influence and are associated with public universities or non-profit civil associations.

In Brazil, there are projects that are focused on an ecosystem [e.g., the Ecomangue project, (Instagram @_ecomangue; the Araçá Bay)] (Amaral et al. 2010, 2015) or a specific group [e.g., Projeto Cephalopoda (Instagram @projeto_cephalopoda)], which are developed through joint action by researchers and local communities. In Uruguay, the environmental education newsletter *Caracol Negro y sus Ovicápsulas* (Scarabino et al. 2020) has been particularly well received and used in a multitude of educational settings. Scarabino (2021) detailed ongoing or particularly interesting activities related to environmental-heritage education focused on molluscs from the country's eastern coast (Rocha). In Argentina, the non-profit entity ProyectoSub promotes ocean conservation, especially that of marine invertebrates and algae, using social networks and audiovisual technologies to encourage oceanic culture (<https://www.proyectosub.org.ar/>). There is also an educational program in Argentina focused on malacology, which contributes to the dissemination of knowledge about molluscs and promotes education in this field (Darrigran et al. 2014).

Management of molluscan fisheries

Molluscan fisheries in the SW Atlantic still lack effective regulatory measures designed to promote the conservation and management of these marine resources.

In Brazil, there is some federal and/or state legislation to regulate the artisanal fishery of the clam *Anomalocardia flexuosa* in some locations (Normative Instruction IBAMA no. 81, 28 December 2005), the artisanal fishery of *Octopus*

insularis in the Fernando de Noronha Archipelago (Leite et al. 2008a, b), the commercial octopus pot fishery in north and northeastern regions (Normative Instruction SEAP/PR no. 15, 19 June 2007), the pot fishery of *Octopus americanus* in the southern region (Normative Instruction SEAP/PR no. 26, of 19 December 2008 and Ordinance SAP/MAPA no. 452, of 18 November 2021), and the squid fishery in Rio de Janeiro (law no. 282–88 and Ordinance ICMBio no. 6, of 29 January 2014) and Santa Catarina (Ordinance SAQ no. 1/2024, of 9 February 2024) states. However, all of these measures are spatially restricted and fail to consider the geographic distribution of the target species. The management of *A. flexuosa*, for example, needs to take into account the structure of subpopulations (banks) at different levels of connectivity within a metapopulation, exploitation in specific banks, and the establishment of protected areas to maintain stocks (Rocha 2013). In the case of *O. insularis*, while the abundance of the Fernando de Noronha Archipelago population appears to be stable due an adequate management plan (Leite et al. 2008a, b), this species has also been artisanally harvested along the tropical Brazilian coast without regulation or monitoring, despite the high number of scientific publications on the species that can support appropriate legislation for its management (Leite et al. 2008a, 2016; Lima et al. 2014a, b). In turn, the change in the minimum depth allowed for capturing *O. americanus* (from 70 m to 35 m deep) (Ordinance SAP/MAPA n°. 452, of November 18, 2021) did not consider the ecology of the species, leading to the proposal of conservation measures, such as restricting industrial fishing near rocky shores (vital areas for juvenile growth and feeding) and maintaining artisanal freediving fishing for subsistence purposes (Côrtes 2022).

In Uruguay, there are also some management and conservation measures used to regulate fishing activity, such as the prohibition of the fishing of *Adelomelon beckii* and the establishment of minimum landing and commercialization sizes for *Mytilus platensis*, *Amarilladesma mactroides* and *Zidona dufresnii* (Resolution DINARA no. 213/018). In the case of *A. beckii*, the implementation of some specific fishing regulations have been recommended, such as the establishment of closed areas to guarantee the population's reproductive success, or the prohibition of the shell trade along the coast to definitively prevent landings of this species (Carranza et al. 2017). With regard to *A. mactroides*, authorization for its harvesting went through several phases, with the opening or closing of the season, based on the recovery of populations that presented episodes of mass mortality linked to harmful algal blooms and an increase in seawater temperature (Gianelli et al. 2021). These changes have significantly impacted the social-ecological system of the area, with the 14-year closure of its fishing in Uruguay resulting in economic losses and affecting the livelihoods of local communities (reviewed in Defeo et al. 2021). Despite

the passage of time, clam populations have not recovered to their pre-mass mortality levels of abundance, indicating their high sensitivity to warming and limited adaptive capacity.

Argentina has also implemented measures to promote sustainable fisheries management, such as quotas and fishing seasons, but these are spatially restricted. The only octopus fishing management regulation (which is currently not applied), for example, is limited to Rio Negro province (northern Patagonia) and establishes rules for *O. tehuelchus* fishing (BO 4288, Resolution no. 025/2005). Regarding the shortfin squid, *Illex argentinus*, within the Argentine EEZ there are fishing management regulations that establish geographical and temporal limits (Consejo Federal Pesquero 2019, Resolution no. 14/2016). However, an important part of the *I. argentinus* life cycle occurs within the limits of the EEZ and in the adjacent international waters, where there is unregulated fishing by foreign overseas fleets. Overfishing in areas adjacent to the national EEZ is becoming a serious problem (FAO 2019), but the effects of the catches on the fished populations are not yet known (Brunetti et al. 2002; Arkhipkin et al. 2015). For *Doryteuthis gahi*, the Malvinas Islands fishery adopted management measures based on effort limitation (Arkhipkin et al. 2015). Regarding marine gastropod catches in North Atlantic Patagonia, a regulation (Regulation no. 199/2018 of the Secretary of Fisheries of Chubut Province) establishes a minimum harvestable size and fishing season for the species *Odontocymbiola magellanica*, *Adelomelon ancilla*, *Adelomelon beckii*, *Zidona dufresnii* (Volutidae), *Buccinastrum deforme*, *Buccinanops cochlidium* (Nassariidae), *Trophon geversianus* (Muricidae), *Tegula patagonica* (Trochidae), and *Nacella magellanica* (Patellidae) (Cumplido and Bigatti 2020; Cumplido et al. 2022). This was established prior to the massive exploitation of the resource, but only in Chubut Province, despite recommendations of specific fishery regulations for the entire coast of Argentina (Giménez et al. 2005; Arrighetti et al. 2011). With regard to the observed populational decline of *Amarilladesma mactroides*, the Buenos Aires Province Fisheries Directorate and the Municipality of Partido de la Costa have implemented measures prohibiting the extraction of this bivalve (Directive no. 956/96 of the Buenos Aires Province Fisheries Directorate, 1238/96, 1704/96, 1004/01).

Using scientific research together with traditional knowledge is also an important initiative in establishing effective regulatory conservation measures. An experiment with the artisanal pot fishery in northeastern Brazil was developed together with local fishers, who collaborated with the planning, developing and evaluation of the efficiency of pots in capturing *Octopus insularis* (Andrade 2016). Interviews with artisanal octopus fishers in several MPAs allowed the redescription and delimitation of the distribution area of the octopus *Callistoctopus furvus* (Jesus et al. 2021a, b, 2022). Ethnoecological research in the Coroa Vermelha

Environmental Protection Area (Brazil) demonstrated that fishers have knowledge about biological and ecological aspects of the octopuses that are compatible with scientific information about the species, and especially about its behavior and trophic ecology (Martins et al. 2011).

Most marine mollusc stocks that are commercially exploited in the warm-temperate SW Atlantic are fully exploited, overexploited, or depleted (Gianelli and Defeo 2017; Gianelli et al. 2023). In industrial fisheries developed in this area, sustainable management systems include onboard and port-based monitoring of catches, surveys and stock assessments, reference points, limited entry, rotational harvesting, and no-take areas (Defeo and Vasconcellos 2020). Considering that most sessile and sedentary mollusc species, and the typical way in which they are fished, have a strong and persistent spatial structure, spatially explicit management considerations should be of utmost priority for them (Defeo and Castilla 2005; Orensanz et al. 2005). For example, the distribution area of the Patagonian scallop is divided into management units, and the stock status of each unit is determined by a combination of biomass estimates and the proportion of harvestable scallops (Soria et al. 2016). Similar spatially explicit considerations have been implemented to manage other relevant commercial bivalve industrial fisheries in the SW Atlantic (Gutiérrez et al. 2011; Masello and Defeo 2016), and have also been considered a relevant management strategy in some small-scale fisheries in the region (Defeo and Castilla 2012; Orensanz et al. 2013).

Challenges and perspectives

Management of marine areas and fisheries

One of the greatest challenges for the effective conservation of marine species is understanding the dynamics of evolutionary and ecological processes in their environment. Consequently, a more nuanced evaluation of extinction risks is necessary, coupled with the implementation of measures for the effective mitigation of oceanic defaunation. The establishment of protected areas and precautionary management of ocean resources can contribute to achieving this objective (McCauley et al. 2015; del Monte-Luna et al. 2023). There are ongoing projects focused on creating new MPAs in the SW Atlantic. In Brazil, there are proposals to establish new MPAs along the coast that have different conservation priorities, and to create a protected area for benthic organisms, including *Octopus americanus*, in the southern region (Magris et al. 2021; Côrtes 2022). The government of Uruguay has made a commitment to increasing the protected marine area of the country to achieve a 30% target by 2030, which will be achieved by creating a network of



Fig. 3 Challenges and prospects for achieving the effective conservation of estuarine and marine mollusc species of the SW Atlantic Ocean

well-connected and representative MPAs (Limongi et al. 2023). In Argentina, marine areas have increased significantly in the last 10 years, with the creation of the Namuncurá Banco Burwood and Yaganes MPAs. The criteria adopted in their creation were based on fisheries and touristic activities, with the aim of protecting economic resources and/or breeding sanctuaries, as well as the protection of marine mammals and birds (Fig. 3).

In the management of the molluscan resources in the SW Atlantic, the prevalence of open access regimes and the extensive accessibility of open coasts to unauthorized fishers and/or recreational harvesters have led to management failures. Additionally, distinct fisher idiosyncrasies among countries and the growing influence of external drivers, such as international trade and market globalization, governance, illegal fishing, and climate change, have emerged as major determinants in landings trends and stock statuses (Defeo and Vasconcellos 2020). Measures designed for the recovery of molluscan resources from such situations need to take into consideration spatially explicit management strategies and a set of biosocioeconomic indicators under a precautionary approach and solid governance schemes that must account for the full participation of fishers (Orensanz et al. 2013). Effective management strategies must take into account the fact that most molluscs are structured as metapopulations (Orensanz et al. 2005), implying that the spatial dimension

of processes related to both stages of the life cycle (i.e., dispersive larvae and benthic adults) must be considered when assessing and managing their stocks. Thus, assessment and management of these stocks require the identification of meaningful spatial scales to provide unbiased estimates of abundance and population structure, and to assess key spatial dimensions (population regulation mechanisms and processes, and the associated spatial dynamics of the harvesting process) (Orensanz et al. 2005; McLachlan and Defeo 2018). Moreover, the high sensitivity of certain exploited species and an increase in the representation of species with potential distribution changes due to climate change pose risks for future catches and benthic fisheries as a whole (Gianelli et al. 2023). Given this complex and concerning situation, comprehensive studies on the ecological impacts of the different external drivers acting jointly, or in isolation, are needed. Additionally, the development of long-term research programs that include the collection of scientific and traditional knowledge, along with participatory monitoring, control, and surveillance of management strategies, could provide a more optimistic perspective than the current one.

Weak governance and institutional instability also pose significant challenges to the sustainability of mollusc fisheries in the SW Atlantic, with important conservation implications. To ensure the long-term sustainability of fisheries, it is crucial to have sustained political commitment, state policies and high-level policy goals, as well as the implementation of ecosystem frameworks and technical management measures. Additionally, fishers themselves should be involved as stewards of their own activities. Incentive-based approaches that clearly define community and individual harvest or territorial rights should be implemented, alongside community-based data collection programs for research, monitoring, and effective oversight, to promote sustainable fisheries and better engage stakeholders in responsible fishing practices (Orensanz et al. 2013). An ecosystem approach to fisheries that includes co-management as a formal mode of governance is suggested to correct failures in fisheries (Gianelli et al. 2018). This approach may be applied, for example, to artisanal, small-scale molluscs fisheries, which are the most common form of exploitation on sandy beaches and rocky shores, and have important socioeconomic relevance in the SW Atlantic, providing food and employment and generating direct incomes for fisher communities and, in some cases, high export revenues for the countries concerned (Defeo et al. 2021). Therefore, management plans are needed to sustain fisheries over time and to contribute significantly to food security and adequate nutrition.

Long-term interdisciplinary monitoring programs

The development of long-term projects can help to establish new protected areas and manage species and their habitats. Although some seemingly reasonable initiatives have been developed in Brazil (Bernardino et al. 2016; Cordeiro et al. 2022), there is still a lack of long-term coordinated studies on molluscs from the SW Atlantic, which is a significant limitation for studying changes in these species and related marine ecosystems over time. For instance, little is known about the ecological impacts on molluscs of the mining dam collapse in 2015 (Matthews-Cascon et al. 2018) and the extensive oil spill in 2019 that affected Brazil's coastline (Viana et al. 2022; Soares and Rabelo 2023). There is also a lack of spatial and temporal trend data that show whether there are site differences and whether levels of pollution are increasing, decreasing, or are stable. The development of long-term monitoring programs that involve the collaboration of multiple countries, organizations and researchers is urgently needed and indispensable. Such programs should use standardized methods for data collection and analysis and involve new technologies, such as DNA sequencing and stable isotope analysis. Furthermore, the use of open access global geospatial datasets could help in delimiting target areas while simultaneously assessing the effects of land use changes (e.g., sediment runoff) in their vicinity (Delevaux and Stamoulis 2022).

Along with long-term monitoring programs, biogeographical studies should also be conducted to promote a better understanding of the biodiversity patterns of SW Atlantic molluscs. In recent years, efforts have been made to understand these diversity patterns in shallow (Barroso et al. 2016a; Cord et al. 2022; Oliveira et al. 2024) and deep waters (Teso et al. 2019; Pacheco et al. 2022), including those of oceanic islands (Hachich et al. 2015; Barroso et al. 2016b). Investigations using this approach can contribute to knowledge of species distribution patterns in different biogeographic areas of this region, and provide insights into the factors that influence benthic patterns on a global scale. Pending challenges in taxonomic and biogeographic research will be successfully addressed only if multinational collaborative initiatives are undertaken in a framework of integrative taxonomy (Scarabino et al. 2015).

Important tools to aid these studies are the on-line databases (WoRMS, MolluscaBase, Obis, GBIF, Malacolog, among others) that contribute to the updating of the taxonomic status and the current distribution range of species, showing and comparing the distribution ranges on a large scale and global level. Thus, the contribution of local and regional editors for the online databases is essential, to clarify the taxonomic status of mollusc species, as is the case with Project Cephalopoda and Lamece—UFSC (Brazil), which are working with photos and video databases

to identify octopus species (O'Brien et al. 2021). Another potential tool is to encourage citizen science efforts for the collection of data on mollusc populations, using platforms such as iNaturalist. This approach has been recognized as a complementary tool in alien species management and large-scale biodiversity conservation (Kelly et al. 2020). Citizen science initiatives can be an effective way of gathering large amounts of data over a long period of time, while also engaging society in reliable research.

Understanding the reproductive characteristics of a species is also a relevant tool that can assist in the creation of MPAs, the management of fisheries, long-term monitoring programs, and biogeographical studies, since the geographical distribution of species is affected, among other factors, by the type of larval development (Barroso et al. 2022). The reproductive characteristics of most mollusc species in the SW Atlantic, including those inhabiting adjacent deep-sea areas, are still little known. However, recent studies have expanded our understanding of deep-sea ecosystems in the area and allowed for the reassessment of species diversity (Pastorino and Scarabino 2008; Penchaszadeh et al. 2016, 2017, 2019, 2022; Averbuj et al. 2018a; Sánchez et al. 2018; Teso and Penchaszadeh 2019; Teso et al. 2019, 2020). These works revealed that the biodiversity and reproductive strategies of various taxa in these extreme conditions are more complex than previously thought. The knowledge gained from these studies is valuable for understanding life in the deep sea and can aid decision-making processes aimed at protecting marine biodiversity.

Major investment by research funding agencies that are disassociated from companies that aim to profit from the exploitation of living and non-living resources, as is the case in Brazil, is especially important for the operation of vessels to study mesophotic and deep-water molluscs. In Brazil, several projects have been developed through different types of cooperation, including between the Brazilian government and navy, research funding agencies and the state-owned Brazilian multinational corporation of the petroleum industry (Petrobras), in addition to partnerships with foreign institutions (see, for example, Absalão 2010; Perez et al. 2012; Veloso et al. 2017; Lavrado et al. 2017). In the case of Argentina, the Puerto Deseado research vessel, which was used in expeditions between 2009 and 2018, is currently out of service for researchers, while public policies to overcome this situation are not evident. An uneven capacity for sampling has also been the case in Uruguay, which mostly relies on international cooperation to achieve an understanding of its deep-sea biodiversity.

Another challenge that needs to be faced by researchers in South America concerns museums and scientific collections, which are fundamental resources for decision-makers (Johnson et al. 2023). Since the study of molluscs in the SW Atlantic began with the European and North

American expeditions of the nineteenth century, most of the specimens collected were deposited in collections in the northern hemisphere. Despite the increase in the number of described species in the last years, many type specimens are housed in European or North American institutions, and access to these materials is often limited. The mismanagement of institutions has also led to the loss of important collections, such as the fire in 2018 at the National Museum of Rio de Janeiro (Zamudio et al. 2018), which housed one of the most important malacological collections in Brazil (Mega 2019). It is imperative that governments and civil society recognize species as heritage and appreciate them for their intrinsic value. Additionally, there is a need for increased investment in the training of taxonomists and other researchers specialized in the study of biodiversity, as well as the development of additional relevant infrastructure. Currently, the number of taxonomists dedicated to the study of marine molluscs is small (approximately 40 Brazilian, 14 Argentine and three Uruguayan researchers), when taking into consideration the vast area and diversity of ecosystems in the SW Atlantic.

Education and scientific dissemination

Effective conservation of marine molluscs requires a multi-faceted approach, including environmental education initiatives to raise public awareness on this issue. Interventions must be carried out in partnership with competent environmental agencies. This integrated and collaborative approach could foster the development of management and conservation strategies for the protection of molluscs and their ecosystems for future generations. The implementation of local marine stewardship initiatives and ocean defenders has been at the forefront of sustainability efforts, but still needs to receive recognition and support (Bennett et al. 2022). In this setting, the Escazu Agreement, developed in Latin America and adopted in March 2018, is the first international treaty in the world to include provisions on the rights of environmental defenders, providing full public access to information, decision-making, and legal protection and recourse concerning environmental matters (United Nations 2018). It also recognizes the right of current and future generations to a healthy environment and sustainable development. The Escazu agreement recognizes the primary role of civil society in securing environmental protection and consequently biodiversity conservation (McLachlan and Defeo 2023). Given their characteristics and socioecological significance across estuarine and marine ecosystems, molluscs possess the potential to be recognized and appreciated as pivotal components within these initiatives.

Conclusions

Despite the abundance of available information on estuarine and marine molluscs in the SW Atlantic, there remains a considerable way to go for their effective conservation. The numerous knowledge gaps highlight the necessity for effective public measures aimed at expanding financial and structural resources to attain the requisite knowledge and apply it more effectively in the establishment of environmental protection areas and for fishery stocks management. One of the pivotal considerations in forming these protected areas is to recognize benthic organisms as an essential component for the conservation of their entire ecosystem. Given the aforementioned, the following crucial measures can be outlined for the enhanced conservation of estuarine and marine molluscs in the SW Atlantic:

1. Strengthening management and governance to conserve habitats and control pollution and non-native species, with the aim of reducing the deterioration of ecosystem services provided by molluscs.
2. Establishing the necessary foundations for the monitoring of environmental changes—particularly those related to climate change and excessive human alterations that could lead to the loss of biodiversity in estuarine and marine ecosystems—through long-term interdisciplinary projects.
3. Expanding and supporting the training of researchers specializing in molluscs.
4. Increasing funding for scientific dissemination projects to increase society's awareness of the importance of molluscs.
5. Enhancing and improving legislation aimed at regulating mollusc fisheries.

Each of these measures requires a strategic approach and collaboration among various stakeholders, including governments, research institutions, local communities, and non-governmental organizations. Successful mollusc conservation depends on both the efficient implementation of short-term projects and investment in long-term solutions to address climate change and ecosystem degradation. Among the short-term projects, whose results could be observed within 2–3 years, are investments in researcher training, with an increase in capacity-building programs for taxonomists and greater funding for scientific dissemination. However, these measures will not be effective without long-term programs focused on integrated coastal zone management, which include partnerships between governments and local communities to regulate land use, pollution, and fishing and shellfish harvesting, as well as the establishment of environmental monitoring networks. Such projects are critically important for assessing real-time climatic and anthropogenic impacts and require planning, legislation, and continuous implementation.

Acknowledgements We are thankful to all of the researchers who have dedicated themselves to malacology. We are also thankful to Alexandre Dias Pimenta for contributing to the early stages of the manuscript, and Comisión Sectorial de Investigación Científica (CSIC Grupos 32).

Author contributions All of the authors contributed to the study conception and design. Material preparation, data collection and analysis were performed by CXB, EPA and VT. The first draft of the manuscript was written by CXB and all of the authors commented on previous versions of the manuscript. All of the authors read and approved the final manuscript.

Funding FDP has received financial support from the São Paulo Research Foundation (FAPESP) (grant no. 2018/10313-0).

Data availability No datasets were generated or analyzed during the current study.

Declarations

Conflict of interest The authors declare no competing interests.

References

- Ab'Saber AN (2005) Brazilian coast. Metalivros, São Paulo
- Absalão RS (2010) Mollusca. In: Lavrado HP, Brasil ACS (eds) Biodiversidade da região oceânica profunda da Bacia de Campos: Macrofauna. SAG Serv, Rio de Janeiro, pp 31–94
- Agnew DJ, Hill SL, Beddington JR, Purchase LV, Wakeford RC (2005) Sustainability and management of southwest Atlantic squid fisheries. *Bull Mar Sci* 76:579–594
- Alves J, Haimovici M (2011) Reproductive biology of *Octopus tehuelchus* d'Orbigny, 1834 (Cephalopoda: Octopodidae) in southern Brazil. *Nautilus* 125:150–158
- Amaral ACZ, Migotto AE, Turra A, Schaeffer-Novelli Y (2010) Araçá: biodiversidade, impactos e ameaças. *Biota Neotrop* 10:219–264. <https://doi.org/10.1590/S1676-06032010000100022>
- Amaral ACZ, Turra A, Ciotti AM, Wongtschowski CLBR, Schaeffer-Novelli Y (2015) Vida na Baía do Araçá: diversidade e importância. Lume, São Paulo
- Amaral ACZ, Corte GNC, Rosa-Filho JS et al (2016) Brazilian sandy beaches: characteristics, ecosystem services, impacts, knowledge and priorities. *Braz J Oceanogr* 64:5–16. <https://doi.org/10.1590/S1679-875920160933064sp2>
- Amaral VS, Simone LRL, Tâmega FTS et al (2020) New records of the non-indigenous oyster *Saccostrea cucullata* (Bivalvia: Ostreidae) from the southeast and south Brazilian coast. *Reg Stud Mar Sci* 33:100924. <https://doi.org/10.1016/j.rsma.2019.100924>
- Andrade MP, Santos CMD, De Paiva MMM et al (2023) Assessing negative welfare measures for wild invertebrates: the case for octopuses. *Animals* 13:3021. <https://doi.org/10.3390/ani13193021>
- Andrade LCA (2016) Estratégias de exploração e comércio da pesca artesanal de polvo. Dissertation, Universidade Federal do Rio Grande do Norte
- Appeltans W, Ah Yong ST, Anderson G et al (2012) The magnitude of global marine species diversity. *Curr Biol* 22:2189–2202. <https://doi.org/10.1016/j.cub.2012.09.036>
- Araújo FC, Maia RC (2021) O uso do gastrópode *Melampus coffea* (Ellobiidae) (Linnaeus, 1758) como indicador ecológico do estado de degradação ambiental de manguezais. *J Integr Coast Zone Manag* 21:25–32. <https://doi.org/10.5894/rgci-n366>

- Arias AH, Vazquez-Botello A, Tombesi N et al (2010) Presence, distribution, and origins of polycyclic aromatic hydrocarbons (PAHs) in sediments from Bahia Blanca estuary, Argentina. *Environ Monit Assess* 160:301–314. <https://doi.org/10.1007/s10661-008-0696-5>
- Arkhipkin AI, Rodhouse PGK, Pierce GJ et al (2015) World squid fisheries. *Rev Fish Sci Aquac* 23:92–252. <https://doi.org/10.1080/23308249.2015.1026226>
- Arrighetti F, Penchaszadeh PE (2010) Gametogenesis, seasonal reproduction and imposex of *Adelomelon beckii* (Neogastropoda: Volutidae) in Mar del Plata, Argentina. *Aquat Biol* 9:63–75. <https://doi.org/10.3354/ab00226>
- Arrighetti F, Brey T, Mackensen A, Penchaszadeh PE (2011) Age, growth and mortality in the giant snail *Adelomelon beckii* (Broderip 1836) on the Argentinean Shelf. *J Sea Res* 65:219–223. <https://doi.org/10.1016/j.seares.2010.10.003>
- Arrighetti F, Landro SM, Lambre ME, Penchaszadeh PE, Teso V (2019) Multiple-biomarker approach in the assessment of the health status of a novel sentinel mussel *Brachidontes rodriguezii* in a harbor area. *Mar Pollut Bull* 140:451–461. <https://doi.org/10.1016/j.marpolbul.2019.02.002>
- Avendaño O, Roura Á, Cedillo-Robles CE et al (2020) *Octopus americanus*: a cryptic species of the *O. vulgaris* species complex re-described from the Caribbean. *Aquat Ecol* 54:909–925. <https://doi.org/10.1007/s10452-020-09778-6>
- Averbuj A, Penchaszadeh PE (2010) On the reproductive biology and impact of imposex in a population of *Buccinanops monilifer* from Mar del Plata, Argentina. *J Mar Biol Assoc U K* 90:729–734. <https://doi.org/10.1017/S0025315409990956>
- Averbuj A, Penchaszadeh PE, Pastorino G (2018a) Egg masses and development of *Falsilunatia eltanini* (Mollusca: Gastropoda): a deep-sea naticid from a southwestern Atlantic canyon. *Mar Biol* 165:81. <https://doi.org/10.1007/s00227-018-3337-8>
- Averbuj A, Primost MA, Giulianelli S, Bigatti G (2018b) Acute toxicity of tributyltin to encapsulated embryos of a marine gastropod. *Molluscan Res* 38:29–33. <https://doi.org/10.1080/13235818.2017.1357671>
- Ávila-da-Silva AO, Assunção R, Tomás ARG (2014) Capítulo 9: Surgimento e evolução da pesca do polvo-comum, *Octopus vulgaris* Cuvier, 1797, com potes do estado de São Paulo, Brasil. In: (eds) Haimovici M, Andriquetto Filho JM, and Sunye OS. A pesca marinha e estuarina no Brasil: estudos de caso multidisciplinares. Editora da FURG, Rio Grande, pp 101–110
- Azevedo D, Rocha-Barreira CA, Matthews-Cascon H, Castro IB (2012) *Pugilina morio* L., a new imposex exhibitor from South American estuarine environments: approach for a non-lethal method to evaluate imposex. *Bull Environ Contam Toxicol* 89:786–792. <https://doi.org/10.1007/s00128-012-0779-4>
- Banha TNS, Luiz OJ, Asp NE et al (2022) The Great Amazon Reef System: a fact. *Front Mar Sci*. <https://doi.org/10.3389/fmars.2022.1088956>
- Barbosa IS, Brito GB, Santos GL et al (2019) Multivariate data analysis of trace elements in bivalve molluscs: characterization and food safety evaluation. *Food Chem* 273:64–70. <https://doi.org/10.1016/j.foodchem.2018.02.063>
- Barletta M, Lima ARA, Costa MF (2019) Distribution, sources and consequences of nutrients, persistent organic pollutants, metals and microplastics in South American estuaries. *Sci Total Environ* 651:1199–1218. <https://doi.org/10.1016/j.scitotenv.2018.09.276>
- Barón PJ, Ré ME (2002) Reproductive cycle and population structure of *Loligo sanpaulensis* of the northeastern coast of Patagonia. *Bull Mar Sci* 71:175–186
- Barroso CX, Lotufo TMC, Matthews-Cascon H (2016a) Biogeography of Brazilian Prosobranch gastropods and their Atlantic relationships. *J Biogeogr* 43:2477–2488. <https://doi.org/10.1111/jbi.12821>
- Barroso CX, Lotufo TMC, Bezerra LEA, Matthews-Cascon H (2016b) A biogeographic approach to the insular marine “Prosobranch” gastropods from the southwestern Atlantic Ocean. *J Molluscan Stud* 82:558–563. <https://doi.org/10.1093/mollus/eyw015>
- Barroso CX, Freitas JEP, Sportorno P et al (2021) Molluscs from the upper mesophotic zone in a scarcely known reef of the western equatorial Atlantic. *Arq Cienc Mar* 54:44–58. <https://doi.org/10.32360/acmar.v54i2.61276>
- Barroso CX, Lotufo TMC, Matos AS, Carneiro PBM, Matthews-Cascon H (2022) The distribution of marine gastropods is more influenced by larval development than by adult characteristics. *Mar Biol* 169:83. <https://doi.org/10.1007/s00227-022-04069-0>
- Bastos P, Vieira GC, Reis IMM, Costa RL, Lopes GR (2018) Feeding behavior of octopus *Octopus vulgaris* type II (Cuvier, 1797) paralarvae fed on *Artemia* enriched with microalgae and supplemented with DHA. *Arq Bras Med Vet Zootec* 70:628–632. <https://doi.org/10.1590/1678-4162-9731>
- Bastos P, Fracalossi DM, Chimal ME, Sánchez A, Rosas C (2020) Digestive enzymes and timing of digestion in *Octopus vulgaris* type II. *Aquac Rep* 16:100262. <https://doi.org/10.1016/j.aqrep.2019.100262>
- Battini N, Giachetti CB, Castro KL, Bortolus A, Schwindt E (2021) Predator-prey interactions as key drivers for the invasion success of a potentially neurotoxic sea slug. *Biol Invasions* 23:1207–1229. <https://doi.org/10.1007/s10530-020-02431-1>
- Beck MW, Brumbaugh RD, Airoldi L et al (2011) Oyster reefs at risk and recommendations for conservation, restoration, and management. *BioSci* 61:107–116. <https://doi.org/10.1525/bio.2011.61.2.5>
- Bennett NJ, Le Billon P, Belhabib D, Satizábal P (2022) Local marine stewardship and ocean defenders. *NPJ Ocean Sustain* 1:3. <https://doi.org/10.1038/s44183-022-00002-6>
- Bernardino AF, Pagliosa PR, Christofoletti RA et al (2016) Benthic estuarine communities in Brazil: moving forward to long-term studies to assess climate change impacts. *Braz J Oceanogr* 64:81–96. <https://doi.org/10.1590/S1679-875920160849064sp2>
- Berrueta M, Desiderio J, Agliano F et al (2020) Mating behaviour of Patagonian octopus (*Octopus tehuelchus*) under laboratory conditions. *Mar Fish Sci* 33(1):115–120. <https://doi.org/10.47193/mafis.3312020061807>
- Beyer J, Green NW, Brooks S et al (2017) Blue mussels (*Mytilus edulis* spp.) as sentinel organisms in coastal pollution monitoring: a review. *Mar Environ Res* 130:338–365. <https://doi.org/10.1016/j.marenvres.2017.07.024>
- Bigatti G, Ciocco NF (2008) Volutid snails as an alternative resource for artisanal fisheries in northern Patagonic gulfs: availability and first suggestions for diving catches. *J Shellfish Res* 27:417–421. [https://doi.org/10.2983/0730-8000\(2008\)27\[417:VSAAAR\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2008)27[417:VSAAAR]2.0.CO;2)
- Bigatti G, Penchaszadeh PE (2005) Imposex in *Odontocymbiola magellanica* (Caenogastropoda: Volutidae) in Patagonia. *Comun Soc Malacol Urug* 9:371–375
- Bigatti G, Signorelli J (2018) Marine invertebrate biodiversity from the Argentine Sea, south western Atlantic. *Zookeys* 791:47–70. <https://doi.org/10.3897/zookeys.791.22587>
- Bigatti G, Primost MA, Cledón M et al (2009) Biomonitoring of TBT contamination and imposex incidence along 4700 km of Argentinean shoreline (SW Atlantic: From 38S to 54S). *Mar Pollut Bull* 58:695–701. <https://doi.org/10.1016/j.marpolbul.2009.01.001>
- Bigatti G, Diaz De Vivar ME, Cumplido M et al (2018) Fatty acids and contaminants in edible marine gastropods from Patagonia. *J Mar Biol Assoc U K* 98:1355–1363. <https://doi.org/10.1017/S0025315417000625>
- Bigatti G, Cumplido M, Averbuj A (2015) Gasterópodos de interés comercial en la Provincia del Chubut. Informe para la Mesa Técnica Zona 1: Reglamentación de la pesca de gasterópodos.

- LAPEMAR, Laboratorio de Peces y Mariscos de Interés Comercial (CENPAT), Chubut, Argentina
- Bocco G, Cinti A, Vezub J, Sánchez-Carnero N, Chávez M (2019) Lugar y sentido de lugar en un camino de la costa atlántica patagónica, 1950–1970. *Reg Soc* 31:e1127. <https://doi.org/10.22198/rys2019/3/1/1127>
- Bökenhans V, Fernández Alfaya JE, Bigatti G, Averbuj A (2019) Diet of the invasive sea slug *Pleurobranchaea maculata* in Patagonian coastal waters. *N Z J Zool* 46:87–94. <https://doi.org/10.1080/03014223.2018.1464035>
- Bökenhans V, Galván D, Bigatti G, Averbuj A (2022) Stable isotopes reveal algal assimilation in the carnivorous sea slug *Pleurobranchaea maculata* (Quoy & Gaimard, 1832) on Patagonian coasts. *Malacologia* 65:1–2. <https://doi.org/10.4002/040.065.0106>
- Bom FC, Colling LA (2022) The bivalves *Amarilladesma mactroides* and *Donax hanleyanus* as bioindicators of the impact of vehicles on Cassino Beach, southern Brazil. *An Acad Bras Cienc* 94:1–11. <https://doi.org/10.1590/0001-3765202202011265>
- Borges FO, Guerreiro M, Santos CP, Paula JR, Rosa R (2022) Projecting future climate change impacts on the distribution of the *Octopus vulgaris* species complex. *Front Mar Sci* 9:1018766. <https://doi.org/10.3389/fmars.2022.1018766>
- Borthagaray AI, Carranza A (2007) Mussels as ecosystem engineers: their contribution to species richness in a rocky littoral community. *Acta Oecol* 31:243–250. <https://doi.org/10.1016/j.actao.2006.10.008>
- Botelho ERO, Santos MC (2005) A cata de crustáceos e moluscos no manguezal do Rio Camaragibe—estado de alagoas: aspectos sócio-ambiental e técnico-econômico. *Bol Tec Cien CEPENE* 13:77–96
- Bouchet P, Bary S, Héros V, Marani G (2016) How many species of molluscs are there in the world's oceans, and who is going to describe them? In: Héros V et al (eds) *Tropical Deep-Sea Benthos* 29. Mémoires du Muséum National d'Histoire Naturelle, Paris, pp 9–24
- Bouchet P, Decock W, Lonnaville B, Vanhoorne B, Vandepitte L (2023) Marine biodiversity discovery: the metrics of new species descriptions. *Front Mar Sci* 10:929989. <https://doi.org/10.3389/fmars.2023.929989>
- Braga R, Crespi-Abril AC, Van der Molen S, Bainy MCRS, Ortiz N (2019) Analysis of the morphological variation of *Doryteuthis sanpaulensis* (Cephalopoda: Loliginidae) in Argentinian and Brazilian coastal waters using geometric morphometrics techniques. *Mar Biodivers* 47:755–762. <https://doi.org/10.1007/s12526-017-0661-z>
- Braga R, Van der Molen S, Pontones J, Ortiz N (2021) Embryonic development, hatching time and newborn juveniles of *Octopus tehuelchus* under two culture temperatures. *Aquaculture* 530:735778. <https://doi.org/10.1016/j.aquaculture.2020.735778>
- Braga R, Van der Molen S, Rodríguez YE et al (2022) Morphophysiological responses of *Octopus tehuelchus* juveniles during the transition period between endogenous and exogenous feeding. *Aquac* 556:738269. <https://doi.org/10.1016/j.aquaculture.2022.738269>
- Breves A, Skinner LF (2014) First record of the vermetid *Petalocochus varians* (d'Orbigny, 1841) on floating marine debris at Ilha Grande, Rio de Janeiro, Brazil. *J Integr Coast Zone Manag* 14:159–161. <https://doi.org/10.5894/rgci457>
- Breves A, Scarabino F, Leoni V (2014) First records of the non-native bivalve *Isognomon bicolor* (C. B. Adams, 1845) rafting to the Uruguayan coast. *Check List* 10:684–686. <https://doi.org/10.15560/10.3.684>
- Breves A, de Paula TS, Spotorno P et al (2022) Living in solitude or building reefs: ecophenotypic variation of the vermetid *Petalocochus varians* revealed by mitochondrial DNA analysis. *J Molluscan Stud* 88:eyac030. <https://doi.org/10.1093/mollus/eyac030>
- Breves-Ramos A, Junqueira AOR, Lavrado HP, Silva SHG, Ferreira-Silva MAG (2010) Population structure of the invasive bivalve *Isognomon bicolor* on rocky shores of Rio de Janeiro State (Brazil). *J Mar Biol Assoc U K* 90:453–459. <https://doi.org/10.1017/S0025315409990919>
- Brunetti NE, Ivanovic ML, Sakai M (1999) Calamares de importancia comercial en la Argentina. Biología, distribución, pesquerías, muestreo biológico. Instituto Nacional de Investigación y Desarrollo Pesquero, Mar del Plata, Argentina
- Brunetti NE, Ivanovic ML, Pineda S (2002) Unidad 4. Pesquerías de invertebrados marinos. I. Calamares. In: JICA-INIDEP, Instituto Nacional de Investigación y Desarrollo Pesquero (eds), Seminario internacional sobre métodos de Evaluación y monitoreo de los recursos pesqueros. INIDEP, Mar del Plata, pp 1–16
- Burone L, Franco-Fraguas P, Carranza A et al (2021) Physical drivers and dominant oceanographic processes on the Uruguayan margin (southwestern Atlantic): a review and a conceptual model. *J Mar Sci Eng* 9:304. <https://doi.org/10.3390/jmse9030304>
- Buzzi NS, Oliva AL, Arias AH, Marcovecchio JE (2017) Assessment of trace metal accumulation in native mussels (*Brachidontes rodriguezii*) from a South American temperate estuary. *Environ Sci Pollut Res* 24:15781–15793. <https://doi.org/10.1007/s11356-017-9237-5>
- Calazans SH, Lourenço CR, Nicastro KR et al (2021) Unlocking the history of a trans-Atlantic invader: Did the human slave trade impact brown mussel dispersal? *J Biogeogr* 48:2671–2681. <https://doi.org/10.1111/jbi.14232>
- Camillo E, Quadros J, Castro ÍB, Fernandez M (2004) Imposex in *Thais rustica* (Mollusca: Neogastropoda) (Lamarck, 1822) as an indicator of organotin compounds pollution at Maceio coast (northeastern Brazil). *Braz J Oceanogr* 52:101–105
- Campagnari B, Geiger DL (2018) How many micromollusks are there? A case study on species richness in Hawaii, with the description of a new species of *Murdochella* (Gastropoda: Epitoniidae). *Nautilus* 132:83–90
- Cardoso RS, Caetano CHS, Cabrini TMB (2009) Biphallia in imposed females of marine gastropods: new record for *Nassarius vibex* from Brazil. *Braz J Biol* 69:223–224. <https://doi.org/10.1590/S1519-69842009000100030>
- Carmona L, Pola M, Gosliner TM, Cervera JL (2017) Integrative taxonomy and biogeography of the genus *Bulbaeolidia* (Nudibranchia: Aeolidida). *J Molluscan Stud* 83:440–450. <https://doi.org/10.1093/mollus/eyx027>
- Carneiro PBM, Ximenes Neto AR, Jucá-Queiroz B et al (2022) Interconnected marine habitats form a single continental-scale reef system in South America. *Sci Rep* 12:17359. <https://doi.org/10.1038/s41598-022-21341-x>
- Carranza A (2006) Large gastropod by-catch in the hake fishery at the Argentinean-Uruguayan common fishing zone. *Comun Soc Malacol Urug* 9:61–67
- Carranza A, Arim M (2023) Assessing the ecological vulnerability of western Atlantic marine benthic gastropods. *Mar Fish Sci* 36:165–178. <https://doi.org/10.47193/mafis.3622023010509>
- Carranza A, Horta S (2008) Megabenthic gastropods in the outer Uruguayan continental shelf: composition, distribution and some effects of trawling. *Rev Biol Mar Oceanogr* 43:137–142. <https://doi.org/10.4067/S0718-19572008000100014>
- Carranza A, Defeo O, Beck M, Castilla JC (2009) Linking fisheries management and conservation in bioengineering species: the case of South American mussels (Mytilidae). *Rev Fish Biol Fish* 19:349–366. <https://doi.org/10.1007/s11600-009-9108-3>
- Carranza A, Mello C, Ligrone A et al (2010) Observations on the invading gastropod *Rapana venosa* in Punta del Este, Maldonado Bay, Uruguay. *Biol Invasions* 12:995–998. <https://doi.org/10.1007/s10530-009-9534-0>

- Carranza A, Defeo O, Gracia A et al (2011) Towards a South American network for shellfish conservation and restoration. *Tentacle* 19:3–10
- Carranza A, Martínez G, Arim M (2017) An endangered giant marine gastropod *Adelomelon beckii* (Broderip, 1836) in Uruguayan waters. *Pan-Am J Aquat Sci* 12(2):117–122
- Carranza A, Agudo-Padrón I, Collado GA et al (2023) Socio-environmental impacts of non-native and transplanted aquatic mollusc species in South America: What do we really know? *Hydrobiologia* 850:1001–1020. <https://doi.org/10.1007/s10750-023-05164-z>
- Carvalho NF, Waters LG, Arantes RCM et al (2023) Underwater surveys reveal deep-sea corals in newly explored regions of the southwest Atlantic. *Comm Earth Environ* 4:282. <https://doi.org/10.1038/s43247-023-00924-0>
- Castro IB, Matthews-Cascon H, Fernandez MA (2000) Imposax em *Thais haemastoma* (Mollusca: Gastropoda): uma indicação da contaminação por organoestânicos na costa do município de Fortaleza—Ceará—Brasil. *Arq Cienc Mar* 33:51–56
- Castro IB, Meirelles CAO, Matthews-Cascon H, Fernandez MA (2004) *Thais (Stramonita) rustica* (Lamarck, 1822) (Mollusca: Gastropoda: Thaididae), a potential bioindicator of contamination by organotin in northeast Brazil. *Braz J Oceanogr* 52:135–139
- Castro IB, Braga ARC, Rocha-Barreira CA (2005a) Altos índices de imposax em *Stramonita rustica* (Mollusca: Gastropoda) em áreas portuárias dos estados de Alagoas e Sergipe, Brasil. *Trop Oceanogr* 33:121–128. <https://doi.org/10.5914/tropocean.v33i2.5056>
- Castro IB, Meirelles CA, Pinheiro JCL, Matthews-Cascon H, Rocha-Barreira CA (2005b) The increasing incidence of imposax in *Stramonita haemastoma* (Mollusca: Gastropoda: Muricidae) after the establishment of the Pecém Harbor, Ceará State, northeast Brazil. *Thalassas* 21:71–75
- Castro IB, Lima AFA, Braga ARC, Rocha-Barreira CA (2007a) Imposax in two muricid species (Mollusca: Gastropoda) from the northeastern Brazilian coast. *J Braz Soc Ecotox* 2:81–91
- Castro IB, Bemvenuti CE, Fillmann G (2007b) Preliminary appraisal of imposax in areas under the influence of southern Brazilian harbors. *J Braz Soc Ecotox* 2:73–79
- Castro IB, Ribeiro-Ferreira VP, Lima AFA, Meirelles CA, Rocha-Barreira CA (2007c) Imposax in three Prosobranch species from Ilha do Japonês, Rio de Janeiro, southeast Brazil. *Thalassas* 23:37–42
- Castro IB, Meirelles CAO, Matthews-Cascon H et al (2008) Imposax in endemic volutid from northeast Brazil (Mollusca: Gastropoda). *Braz Arch Biol Technol* 51:1065–1069. <https://doi.org/10.1590/S1516-89132008000500024>
- Castro IB, Rossato M, Fillmann G (2012) Imposax reduction and residual butyltin contamination in southern Brazilian harbors. *Environ Toxicol Chem* 31:947–954. <https://doi.org/10.1002/etc.1793>
- Cavaleiro NP, Lazoski C, Tureck CR et al (2019) *Crassostrea talonata*, a new threat to native oyster (Bivalvia: Ostreidae) culture in the southwest Atlantic. *J Exp Mar Bio Ecol* 511:91–99. <https://doi.org/10.1016/j.jembe.2018.11.011>
- Cetra N, Roche A (2023) Nudibranchia (Mollusca: Gastropoda) from San Matías Gulf, northern Argentine Patagonia. *Zootaxa* 5244(5):455–473. <https://doi.org/10.11646/zootaxa.5244.5.3>
- Christo SW, Staichak G, Vidolin D, Ferreira AL (2021) Preliminary data indicates the importance of depuration in oysters for microfibers contamination. *Pan-Am J Aquat Sci* 16:255–260. <https://doi.org/10.54451/PanamJAS.16.3.255>
- Cinti A, Soria G, Orensanz JM, Parma AM (2003) Relevamiento del sector pesquero artesanal y deportivo en el área del Polo Pesquero Bahía Camarones, provincia del Chubut. Technical report no. 8. Comisión Técnica Dirección de Pesca-Centro Nacional Patagónico (CENPAT), Asociación de Pescadores Artesanales de Puerto Madryn (APAPM), Argentina
- Claremont M, Williams ST, Barraclough TG, Reid DG (2011) The geographic scale of speciation in a marine snail with high dispersal potential. *J Biogeogr* 38:1016–1032. <https://doi.org/10.1111/j.1365-2699.2011.02482.x>
- Cledón M, Theobald N, Gerwinsky W, Penchaszadeh PE (2006) Imposax and organotin compounds in marine gastropods and sediments from the Mar del Plata coast, Argentina. *J Mar Biol Assoc U K* 86:751–755. <https://doi.org/10.1017/S002531540601366X>
- Cledón M, Tremblay LA, Griffiths C et al (2021) Trace metal residues in marine mussels: a global survey. *Environ Toxicol Chem* 40:3434–3440. <https://doi.org/10.1002/etc.5228>
- Commendatore MG, Franco MA, Costa PG et al (2015) Butyltins, polyaromatic hydrocarbons, organochlorine pesticides, and polychlorinated biphenyls in sediments and bivalve mollusks in a mid-latitude environment from the Patagonian coastal zone. *Environ Toxicol Chem* 34:2750–2763. <https://doi.org/10.1002/etc.3134>
- Comoglio L, Amin O, Botté S, Marcovecchio J (2011) Use of biomarkers in resident organisms as a tool for environmental monitoring in a cold coastal system, Tierra del Fuego Island. *Ecotoxicol Environ Saf* 74:382–393. <https://doi.org/10.1016/j.ecoenv.2010.10.005>
- Conti ME, Stripeikis J, Finoia MG, Tudino MB (2011) Baseline trace metals in bivalve molluscs from the Beagle Channel, Patagonia (Argentina). *Ecotoxicology* 20:1341–1353. <https://doi.org/10.1007/s10646-011-0690-5>
- Conti ME, Tudino MB, Finoia MG, Simone C, Stripeikis J (2019) Managing complexity of marine ecosystems: from the monitoring breakdown structure (MBS) to the baseline assessment. Trace metal concentrations in biomonitors of the Beagle Channel, Patagonia (2005–2012). *Ecol Indic* 104:296–305. <https://doi.org/10.1016/j.ecolind.2019.05.013>
- Cooley S, Schoeman D, Bopp L et al (2022) Oceans and coastal ecosystems and their services. In: Pörtner HO, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Löschke S, Möller V, Okem A, Rama B (eds) *Climate change 2022: impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, pp 379–550
- Cord I, Nunes LT, Barroso CX et al (2022) Brazilian marine biogeography: a multi-taxa approach for outlining sectorization. *Mar Biol* 169:61. <https://doi.org/10.1007/s00227-022-04045-8>
- Cordeiro CAMM, Aued AW, Barros F et al (2022) Long-term monitoring projects of Brazilian marine and coastal ecosystems. *PeerJ* 10:e14313. <https://doi.org/10.7717/peerj.14313>
- Cordone G, Lozada M, Vilacoba E et al (2022) Metabarcoding, direct stomach observation and stable isotope analysis reveal a highly diverse diet for the invasive green crab in Atlantic Patagonia. *Biol Invasions* 24:505–526. <https://doi.org/10.1007/s10530-021-02659-5>
- Coronato AMJ, Coronato F, Mazzoni E, Vázquez M (2008) The physical geography of Patagonia and Tierra del Fuego. *Develop Quatern Sci* 11:13–55. [https://doi.org/10.1016/S1571-0866\(07\)10003-8](https://doi.org/10.1016/S1571-0866(07)10003-8)
- Corrêa PVF, Miranda MS, Passos FD (2018) South America-Africa missing links revealed by the taxonomy of deep-sea molluscs: examples from prochaetodermatid aplacophorans. *Deep-Sea Res I Oceanogr Res Pap* 132:16–28. <https://doi.org/10.1016/j.dsr.2017.12.008>
- Corte GN, Esmaeili YS, Maria TF et al (2023) The science we need for the beaches we want: frontiers of the flourishing Brazilian ecological sandy beach research. *Front Mar Sci*. <https://doi.org/10.3389/fmars.2023.1200979>
- Côrtes MO (2022) Distribuição e uso de habitat do polvo *Octopus americanus* Montfort, 1802 (Cephalopoda: Octopodidae):

- informações para manejo e conservação. Dissertation, Universidade Federal de Santa Catarina, Florianópolis
- Cortez FS, Souza LDS, Guimarães LL et al (2018) Ecotoxicological effects of losartan on the brown mussel *Perna perna* and its occurrence in seawater from Santos Bay (Brazil). *Sci Total Environ* 637–638:1363–1371. <https://doi.org/10.1016/j.scitotenv.2018.05.069>
- Cortez FS, Souza LS, Guimarães LL et al (2019) Marine contamination and cytogenotoxic effects of fluoxetine in the tropical brown mussel *Perna perna*. *Mar Pollut Bull* 141:366–372. <https://doi.org/10.1016/j.marpolbul.2019.02.065>
- Costa PAS, Haimovici M (1990) A pesca de polvos e lulas no litoral do Rio de Janeiro. *Cienc Cult* 42:1124–1130
- Costa MB, Otegui MBP, Zamprogno GC et al (2022) Abundance, composition, and distribution of microplastics in intertidal sediment and soft tissues of four species of Bivalvia from southeast Brazilian urban beaches. *Sci Total Environ* 857:159352–159412. <https://doi.org/10.1016/j.scitotenv.2022.159352>
- Costello MJ, Vale MM, Kiessling W, Maharaj S, Price J, Talukdar GH (2022) Cross-chapter paper 1. Biodiversity hotspots. In: Pörtner HO, Roberts DC, Tignor M, Poloczanska ES, Mintenbeck K, Alegría A, Craig M, Langsdorf S, Löschke S, Möller V, Okem A, Rama B (eds) *Climate change 2022: impacts, adaptation and vulnerability. Contribution of Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, New York, NY, pp 2123–2161
- Cowie RH, Bouchet P, Fontaine B (2022) The sixth mass extinction: fact, fiction or speculation? *Biol Rev* 97:640–663. <https://doi.org/10.1111/brv.12816>
- Crespi Abril AC, Morsan EM, Barón PJ (2008) Contribution to understanding the population structure and maturation of *Illex argentinus* (Castellanos, 1960): the case of the inner-shelf spawning groups in San Matias Gulf (Patagonia, Argentina). *J Shellfish Res* 27:225–231. <https://doi.org/10.2983/0730-8000-27.5.1225>
- Cumplido M, Bigatti G (2020) Maturity assessment for the implementation of the first fishery regulation in Patagonian marine gastropods. *Malacologia* 63:139. <https://doi.org/10.4002/040.063.0114>
- Cumplido M, Trobbiani G, Carranza A, Bigatti G (2022) Limited gastropod abundances call for selective, small-scale artisanal fisheries in a Patagonian marine protected area. *Fish Res* 250:106291. <https://doi.org/10.1016/j.fishres.2022.106291>
- Dadon JR, Castaño C, Pérez García R, Chiappini MTG, Cruzes JM (2001) Efectos a largo plazo de las pesquerías de almeja amarilla (*Mesodesma macroides*) y berberecho (*Donax hanleyanus*), y la urbanización sobre las comunidades intermareales. In: Alveal K, Antesana T (eds) *Sustentabilidad de la biodiversidade*. Universidad de Concepción, Chile, pp 703–715
- Darrigran G, Pastorino G, Damborenea C (2014) Applied malacology: a different approach teaching a traditional topic. In: Toledo Piza AR, Tallarico L, Orlandi Introíni G, Barbosa dos Santos S (eds) *Eleventh International Congress on Medical and Applied Malacology*. Cambridge Scholars Publishing, pp 47–63
- Darrigran G, Agudo-Padrón I, Baez P et al (2020) Non-native mollusks throughout South America: emergent patterns in an understudied continent. *Biol Invasions* 22:853–871. <https://doi.org/10.1007/s10530-019-02178-4>
- Darrigran G, Belz C, Carranza A, Collado G, Correoso M, Fabres A, Gutiérrez Gregoric DE, Lodeiros C, Pastorino G, Penchaszadeh P, Salvador R, Santos S, Thiengo S, Damborenea C (2025) What do we know about non-native and invasive aquatic molluscs in South America? *Biology* 14:151. <https://doi.org/10.3390/biology14020151>
- De Biasi JB, Tomás ARG, Hilsdorf AWS (2016) Molecular evidence of two cryptic species of *Stramonita* (Mollusca, Muricidae) in the southeastern Atlantic coast of Brazil. *Genet Mol Biol* 39:392–397. <https://doi.org/10.1590/1678-4685-GMB-2015-0199>
- De Leo FC, Bernardino AF, Sumida PYG (2020) Continental slope and submarine canyons: benthic biodiversity and human impacts. In: Sumida PYG, Bernardino AF, De Leo FC (eds) *Brazilian deep-sea biodiversity*. Springer, pp 37–72
- de Oliveira Soares M, da Cruz Lotufo TM, Vieira LM et al (2016) Brazilian marine animal forests: a new world to discover in the southwestern Atlantic. In: Rossi S, Bramanti L, Gori A, Orejas C (eds) *Marine animal forests*, 1st edn. Springer, pp 73–110
- Defeo O, Castilla JC (2005) More than one bag for the world fishery crisis and keys for co-management successes in selected artisanal Latin American shellfisheries. *Rev Fish Biol Fish* 15:265–283. <https://doi.org/10.1007/s11160-005-4865-0>
- Defeo O, Castilla JC (2012) Governance and governability of coastal shellfisheries in Latin America and the Caribbean: multi-scale emerging models and effects of globalization and climate change. *Curr Opin Environ Sustain* 4:344–350. <https://doi.org/10.1016/j.cosust.2012.05.002>
- Defeo O, McLachlan A, Armitage D, Elliott M, Pittman J (2021) Sandy beach social-ecological systems at risk: regime shifts, collapses, and governance challenges. *Front Ecol Environ* 19:564–573. <https://doi.org/10.1002/fee.2406>
- Defeo O, Vasconcellos M (2020) Transición hacia un enfoque ecosistémico de la pesca. *FAO Documento Técnico de Pesca y Acuicultura* 668. FAO, Rome
- Defeo O, Puig P, Horta S, De Álava A (2011) Coastal fisheries of Uruguay. In: Salas S, Chuenpagdee R, Charles A, Seijo JC (eds) *Coastal fisheries of Latin America and the Caribbean*. FAO fisheries and aquaculture technical paper no. 544. FAO, Rome, pp 357–384
- del Monte-Luna P, Nakamura M, Vicente A et al (2023) A review of recent and future marine extinctions. *Cambridge Prisms Extinction* 1:e13. <https://doi.org/10.1017/ext.2023.11>
- Delevaux JMS, Stamoulis KA (2022) Prioritizing forest management actions to benefit marine habitats in data-poor regions. *Conserv Biol* 36:e13792. <https://doi.org/10.1111/cobi.13792>
- Di Luca J, Pastorino G (2018) A revision of the genus *Savatieria* Rochebrune & Mabilie, 1885: an endemic group of buccinulid gastropods from the Magellanic region. *J Molluscan Stud* 84:293–302. <https://doi.org/10.1093/mollus/eyy019>
- Di Salvatore P, Calcagno JA, Ortiz N, Ríos de Molina MC, Sabatini SE (2013) Effect of seasonality on oxidative stress responses and metal accumulation in soft tissues of *Aulacomya atra*, a mussel from the South Atlantic Patagonian coast. *Mar Environ Res* 92:244–252. <https://doi.org/10.1016/j.marenvres.2013.10.004>
- Dias TLP, Leo Neto NA, Alves RRN (2011) Molluscs in the marine curio and souvenir trade in NE Brazil: species composition and implications for their conservation and management. *Biodivers Conserv* 20:2393–2405. <https://doi.org/10.1007/s10531-011-9991-5>
- Dias TLP, Mota ELS, Gondim AI et al (2013) *Isognomon bicolor* (C. B. Adams, 1845) (Mollusca: Bivalvia): first record of this invasive species for the states of Paraíba and Alagoas and new records for other localities of northeastern Brazil. *Check List* 9:157. <https://doi.org/10.15560/9.1.157>
- Dias PJ, Gilg MR, Lukehurst SS et al (2018) Genetic diversity of a hitchhiker and prized food source in the Anthropocene: the Asian green mussel *Perna viridis* (Mollusca, Mytilidae). *Biol Invasions* 20:1749–1770. <https://doi.org/10.1007/s10530-018-1659-6>
- Domaneschi O, Martins CM (2002) *Isognomon bicolor* (C.B. Adams) (Bivalvia, Isognomonidae): primeiro registro para o Brasil, redescricao da espécie e considerações sobre a ocorrência e distribuição de *Isognomon* na costa brasileira. *Rev Bras Zool* 19:611–627. <https://doi.org/10.1590/S0101-81752002000200017>

- Dominguez JML (2006) The coastal zone of Brazil: an overview. *J Coastal Res* 39:16–20
- Dornellas AP, Graboski R, Hellberg ME, Lotufo TMC (2022) Phylogeography of *Agathistoma* (Turbinidae, Tegulinae) snails in tropical and southwestern Atlantic. *Zool Scripta* 51:76–90. <https://doi.org/10.1111/zsc.12517>
- dos Santos H, Bertollo J, Creed J (2023) Range extension of the Asian green mussel *Perna viridis* (Linnaeus, 1758) into a marine extractive reserve in Brazil. *BioInvasions Rec* 12:208–222. <https://doi.org/10.3391/bir.2023.12.1.17>
- Duarte CA, Giarratano E, Amin OA, Comoglio LI (2011) Heavy metal concentrations and biomarkers of oxidative stress in native mussels (*Mytilus edulis chilensis*) from Beagle Channel coast (Tierra del Fuego, Argentina). *Mar Pollut Bull* 62:1895–1904. <https://doi.org/10.1016/j.marpolbul.2011.05.031>
- Elías R, Sabatini SE, Dávila C (2022) The response of the natural and sewage-impacted intertidal mussel community of the SW Atlantic to pulse (before/after summer) and chronic sewage discharges in the 1997–2014 period. *Mar Fish Sci* 35(3):333–371. <https://doi.org/10.47193/mafis.3532022010901>
- Elías I, Carozza C, Di Giacomo EE et al (2011) 2. Coastal fisheries of Argentina. In: Salas S, Chuenpagdee R, Charles A, Seijo JC (eds) Coastal fisheries of Latin America and the Caribbean. FAO fisheries and aquaculture technical paper no. 544, Rome, pp 13–48
- FAO (2019) The state of world fisheries and aquaculture 2019
- FAO (2022) The state of world fisheries and aquaculture 2022
- Fernandes FC, Rapagnã LC, Bueno GBD (2004) Estudo da população do bivalve exótico *Isognomon bicolor* (C. B. Adams, 1845) (Bivalvia, Isognomonidae) na Ponta da Fortaleza em Arraial do Cabo, RJ. In: Silva J, Souza R (eds) Água de lastro e bioinvasão. Interciência, Rio de Janeiro, pp 133–141
- Fernandes MR, Salgueiro F, de Paula TS, Lôbo-Hajdu G, Pimenta AD (2021a) Cryptic speciation in the “*Marshallora nigrocincta*” species complex (Gastropoda, Triphoridae) from the western Atlantic. *J Zool Syst Evol Res* 59:819–838. <https://doi.org/10.1111/jzs.12461>
- Fernandes MR, Souza LS, Pimenta AD (2021b) Taxonomic assessment of Triphoroidea and Eulimidae (Caenogastropoda) from the Rio Grande Rise, southwestern Atlantic Ocean. *Arch Molluskenkd* 150(1):65–77. <https://doi.org/10.1127/arch.moll/150/065-077>
- Fernandez MA, Limaverde AM, Castro IB, Almeida ACM, Wagener ALR (2002) Occurrence of imposex in *Thais haemastoma*: possible evidence of environmental contamination derived from organotin compounds in Rio de Janeiro and Fortaleza. *Brazil Cad Saude Publica* 18:463–476. <https://doi.org/10.1590/S0102-311X2002000200011>
- Fernández Severini MD, Villagran DM, Buzzi NS, Sartor GC (2019) Microplastics in oysters (*Crassostrea gigas*) and water at the Bahía Blanca Estuary (southwestern Atlantic): an emerging issue of global concern. *Reg Stud Mar Sci* 32:100829. <https://doi.org/10.1016/j.rsma.2019.100829>
- Ferreira GVB, Justino AKS, Eduardo LN et al (2022) Plastic in the inferno: microplastic contamination in deep-sea cephalopods (*Vampyroteuthis infernalis* and *Abralia veranyi*) from the southwestern Atlantic. *Mar Pollut Bull* 174:113309. <https://doi.org/10.1016/j.marpolbul.2021.113309>
- Floyd M, Mizuyama M, Obuchi M et al (2020) Functional diversity of reef molluscs along a tropical-to-temperate gradient. *Coral Reefs* 39:1361–1376. <https://doi.org/10.1007/s00338-020-01970-2>
- Fonseca VB, Guerreiro ADS, Vargas MA, Sandrini JZ (2020) Effects of DCOIT (4,5-dichloro-2-octyl-4-isothiazolin-3-one) to the haemocytes of mussels *Perna perna*. *Comp Biochem Physiol C Toxicol Pharmacol* 232:108737. <https://doi.org/10.1016/j.cbpc.2020.108737>
- Fontes MK, Gusso-Choueri PK, Maranhão LA et al (2018) A tiered approach to assess effects of diclofenac on the brown mussel *Perna perna*: a contribution to characterize the hazard. *Water Res* 132:361–370. <https://doi.org/10.1016/j.watres.2017.12.077>
- Francini-Filho RB, Asp NE, Siegle E et al (2018) Perspectives on the Great Amazon Reef: extension, biodiversity, and threats. *Front Mar Sci*. <https://doi.org/10.3389/fmars.2018.00142>
- Francioni E, Wagener ALR, Calixto RC, Bastos GC (2004) Evaluation of *Perna perna* (Linné, 1758) as a tool to monitoring trace metals contamination in estuarine and coastal waters of Rio de Janeiro, Brazil. *J Braz Chem Soc* 15(1):103–110. <https://doi.org/10.1590/S0103-50532004000100016>
- Francioni E, Wagener A, Scofield AL, Cavalier B (2005) Biomonitoring of polycyclic aromatic hydrocarbon in *Perna perna* from Guanabara Bay. *Brazil Environ Forensics* 6(4):361–370. <https://doi.org/10.1080/15275920500351759>
- Francioni E, Wagener ALR, Scofield AL, Depledge MH, Cavalier B (2007a) Evaluation of the mussel *Perna perna* as a biomonitor of polycyclic aromatic hydrocarbon (PAH) exposure and effects. *Mar Pollut Bull* 54:329–338. <https://doi.org/10.1016/j.marpolbul.2006.11.003>
- Francioni E, Wagener ALR, Scofield AL et al (2007b) Polycyclic aromatic hydrocarbon in inter-tidal mussel *Perna perna*: space-time observations, source investigation and genotoxicity. *Sci Total Environ* 372:515–531. <https://doi.org/10.1016/j.scitotenv.2006.08.046>
- Franco BC, Defeo O, Piola AR et al (2020) Climate change impacts on the atmospheric circulation, ocean, and fisheries in the southwest South Atlantic Ocean: a review. *Clim Change* 162:2359–2377. <https://doi.org/10.1007/s10584-020-02783-6>
- Freitas TBN, Leite TS, Ramos B, di Cosmo A, Proietti MC (2022) In an octopus’s garden in the shade: underwater image analysis of litter use by benthic octopuses. *Mar Pollut Bull* 175:113339. <https://doi.org/10.1016/j.marpolbul.2022.113339>
- Freitas TBN, Leite TS, Ramos B, Proietti MC (2023) Understanding the interactions between cephalopods and marine litter: a research evaluation with identification of gaps and future perspectives. *Mar Pollut Bull* 190:114814. <https://doi.org/10.1016/j.marpolbul.2023.114814>
- García-Méndez K, Padula V, Valdés Á (2022) Integrative systematics of the genus *Dondice* Marcus, 1958 (Gastropoda, Nudibranchia, Myrrhinidae) in the western Atlantic. *Mar Biodivers* 52:42. <https://doi.org/10.1007/s12526-022-01273-5>
- Gaurisais DY, Bernardino AF (2023) Benthic biogeographic patterns on the deep Brazilian margin. *PeerJ* 11:e14585. <https://doi.org/10.7717/peerj.14585>
- Gernet MV, Belz CE, Baggio RA et al (2019) *Nassarius foveolatus* (Gastropoda, Nassariidae), a new record of an exotic species in Brazil. *Pap Avulsos Zool* 59:e20195955. <https://doi.org/10.11606/1807-0205/2019.59.55>
- Gianelli I, Defeo O (2017) Uruguayan fisheries under an increasingly globalized scenario: long-term landings and bioeconomic trends. *Fish Res* 190:53–60. <https://doi.org/10.1016/j.fishres.2017.02.002>
- Gianelli I, Horta S, Martínez G, de la Rosa A, Defeo O (2018) Operationalizing an ecosystem approach to small-scale fisheries in developing countries: the case of Uruguay. *Mar Policy* 95:180–188. <https://doi.org/10.1016/j.marpol.2018.03.020>
- Gianelli I, Ortega L, Pittman J, Vasconcelos M, Defeo O (2021) Harnessing scientific and local knowledge to face climate change in small-scale fisheries. *Global Environ Change* 68:102253. <https://doi.org/10.1016/j.gloenvcha.2021.102253>
- Gianelli I, Orlando L, Cardoso LG et al (2023) Sensitivity of fishery resources to climate change in the warm-temperate southwest Atlantic Ocean. *Reg Environ Change* 23:49. <https://doi.org/10.1007/s10113-023-02049-8>
- Giarratano E, Amin OA (2010) Heavy metals monitoring in the southernmost mussel farm of the world (Beagle Channel, Argentina).

- Ecotoxicol Environ Saf 73:1378–1384. <https://doi.org/10.1016/j.ecoenv.2010.06.023>
- Giarratano E, Gil MN, Malanga G (2011) Seasonal and pollution-induced variations in biomarkers of transplanted mussels within the Beagle Channel. *Mar Pollut Bull* 62:1337–1344. <https://doi.org/10.1016/j.marpolbul.2011.03.037>
- Giarratano E, Gil MN, Malanga G (2014) Biomarkers of environmental stress in gills of ribbed mussel *Aulacomya atra atra* (Nuevo Gulf, northern Patagonia). *Ecotoxicol Environ Saf* 107:111–119. <https://doi.org/10.1016/j.ecoenv.2014.05.003>
- Gil MN, Torres A, Harvey M, Esteves JL (2006) Metales pesados en organismos marinos de la zona costera de la Patagonia argentina continental. *Rev Biol Mar Oceanogr* 41:167–176. <https://doi.org/10.4067/S0718-19572006000200004>
- Gil GM, Troncoso JS, Thomé JW (2007) Manual para manejo e otimização da exploração comercial de moluscos bivalves. Edição do autor, Porto Alegre
- Giménez J, Lasta M, Bigatti G, Penchaszadeh PE (2005) Exploitation of the volute snail *Zidona dufresnei* in Argentine waters, southwestern Atlantic Ocean. *J Shellfish Res* 24:1135–1140. [https://doi.org/10.2983/0730-8000\(2005\)24\[1135:EOTVSZ\]2.0.CO;2](https://doi.org/10.2983/0730-8000(2005)24[1135:EOTVSZ]2.0.CO;2)
- Gleadall IG, Guerrero-Kommritz J, Hochberg FG, Laptikhovskiy VV (2010) The inkless octopuses (Cephalopoda: Octopodidae) of the southwest Atlantic. *Zool Sci* 27:528–553. <https://doi.org/10.2108/zsj.27.528>
- Gofas S, Zenetos A (2003) Exotic molluscs in the Mediterranean basin: current status and perspectives. *Oceanogr Mar Biol* 41:237–277
- Goldberg R, Averbuj A, Cledón M, Luzzatto D, Nudelman NS (2004) Search for triorganotin along the Mar del Plata (Argentina) marine coast: finding of tributyltin in egg capsules of a snail *Adelomelon brasiliana* (Lamarck, 1822) population showing imposex effects. *Appl Organomet Chem* 18:117–123. <https://doi.org/10.1002/aoc.590>
- Gomes RS (2010) *Conus jaspideus* (Mollusca: Neogastropoda: Conoidea) on the Brazilian coast. *J Mar Biol Assoc U K* 91:531–538. <https://doi.org/10.1017/s0025315410000093>
- González AF, Rodhouse PG (1998) Fishery biology of the seven star flying squid *Martialia hyadesi* at South Georgia during winter. *Polar Biol* 19:231–236. <https://doi.org/10.1007/s0030000050239>
- Gouveia N, Harayashiki CAY, Márquez F et al (2022a) Mollusc shell shape as pollution biomarkers: Which is the best biological model? *Mar Pollut Bull* 179:113663. <https://doi.org/10.1016/j.marpolbul.2022.113663>
- Gouveia N, Oliveira AJLA, Harayashiki CAY et al (2022b) Chemical contamination in coastal areas alters shape, resistance and composition of carnivorous gastropod shells. *Chemosphere* 307:135926. <https://doi.org/10.1016/j.chemosphere.2022.135926>
- Guerra Á, Portela JM, del Río JL (2011) Cephalopods caught in the outer Patagonian Shelf and its upper and medium slope in relation to the main oceanographic features. *Fish Res* 109:179–186. <https://doi.org/10.1016/j.fishres.2011.02.003>
- Güller M, Liuzzi MG, Zelaya DG (2015) A new species of *Leptochiton* (Polyplacophora: Leptochitonidae) from the southwestern Atlantic. *Malacologia* 58:147–155. <https://doi.org/10.4002/040.058.0203>
- Gutiérrez NL, Masello A, Uscudun G, Defeo O (2011) Spatial distribution patterns in biomass and population structure of the deep sea red crab *Chaceon notialis* in the southwestern Atlantic Ocean. *Fish Res* 110:59–66. <https://doi.org/10.1016/j.fishres.2011.03.012>
- Hachich NF, Bonsall MB, Arraut EM et al (2015) Island biogeography: patterns of marine shallow-water organisms in the Atlantic Ocean. *J Biogeogr* 42:1871–1882. <https://doi.org/10.1111/jbi.12560>
- Haimovici M, Perez JAA (1991) Abundância e distribuição de cefalópodes em cruzeiros de prospecção pesqueira demersal na plataforma externa e talude continental do sul do Brasil. *Atlântica* 13(1):189–200
- Haimovici M, Costa PAS, Santos RA, Martins AS, Olavo G (2007) Composição de espécies, distribuição e abundância de cefalópodes do talude da região central do Brasil. In: Costa PAS, Olavo G, Martins AS (eds) Biodiversidade da fauna marinha profunda na costa central brasileira. Museu Nacional, Rio de Janeiro, pp 109–132
- Haimovici M, Leite TS, Marinho RA, Batista B, Madrid RM, Lins Oliveira JE, Lima FD, Candice L (2014) 13. As pescarias de polvos do nordeste do Brasil. In: Haimovici M, Andriquetto Filho JM, PS Sunye (eds) A pesca marinha e estuarina no Brasil: estudos de caso multidisciplinares. Editora da FURG, Rio Grande, pp 147–160
- Harayashiki CAY, Sadauskas-Henrique H, Souza-Barros LR et al (2021) Shell form and enzymatic alterations in *Lottia subrugosa* (Gastropoda, Lotiidae) transplanted to a contaminated site. *Mar Pollut Bull* 164:112075. <https://doi.org/10.1016/j.marpolbul.2021.112075>
- He Q, Silliman BR (2019) Climate change, human impacts, and coastal ecosystems in the Anthropocene. *Curr Biol* 29:R1021–R1035. <https://doi.org/10.1016/j.cub.2019.08.042>
- Henriques MB, Marques HLA, Pereira OM, Bastos GCC (2004) Aspectos da estrutura populacional do mexilhão *Perna perna*, relacionados à extração em bancos naturais da Baía de Santos, estado de São Paulo, Brasil. *Bol Inst Pesca* 30:117–126
- Horta PA, Riul P, Amado Filho GM et al (2016) Rhodoliths in Brazil: Current knowledge and potential impacts of climate change. *Braz J Oceanogr* 64:117–135. <https://doi.org/10.1590/S1679-875920160870064sp2>
- Horta P, Koerich G, Grimaldi G, Mueller C, Destri G, Carneiro P (2022) Patagonian marine forests in a scenario of global and local stressors. In: Helbling EW, Narvarte MA, González RA, Villafañe VE (eds) Global change in Atlantic coastal patagonian ecosystems, natural and social sciences of Patagonia. Springer, pp 151–176
- Instituto Internacional Arayara (2023) Análises do leilão de petróleo e gás: diagnóstico do risco socioambiental do 4o ciclo da oerta permanente da ANP. Instituto Internacional Arayara, Brasília, DF
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) (2019) Global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (version 1). IPBES secretariat, Bonn, Germany
- Iribarne OO (1990) Use of shelter by the small Patagonian octopus *Octopus tehuelchus*: availability, selection and effects on fecundity. *Mar Ecol Prog Ser* 66:251–258. <https://doi.org/10.3354/meps066251>
- Ivanov DL, Scheltema AH (2008) Western Atlantic Prochaetodermitidae from 35°N south to the Argentine Basin including the Gulf of Mexico (Mollusca: Aplacophora). *Zootaxa* 1885:1–60. <https://doi.org/10.11646/zootaxa.1885.1.1>
- Jardim JA, Almeida SM, Simone LRL (2022) Taxonomic study on the Polyplacophora (Chitonida: Ischnochitonidae et Acanthochitonidae) collected by the Marion Dufresne (MD55) expedition, with description of a new species. *Zoosystema* 44:151–157. <https://doi.org/10.5252/zoosystema2022v44a5>
- Jesus MD, Sales JBL, Martins RS et al (2021a) Traditional knowledge aids description when resolving the taxonomic status of unsettled species using classical and molecular taxonomy: the case of the shallow-water octopus *Callistoctopus furvus* (Gould, 1852) from the western Atlantic Ocean. *Front Mar Sci* 7:595244. <https://doi.org/10.3389/fmars.2020.595244>
- Jesus MD, Zapelini C, Schiavetti A (2021b) Can citizen science help delimit the geographical distribution of a species? The case of the *Callistoctopus* sp. (“eastern octopus”) on the Brazilian

- coast. *Ethnobiol Conserv.* <https://doi.org/10.15451/ec2020-09-10.03-1-15>
- Jesus MD, Zapelini C, Santana RO, Schiavetti A (2022) Octopus fishing and new information on ecology and fishing of the shallow-water octopus *Callistoctopus furvus* (Gould, 1852) based on the local ecological knowledge of octopus fishers in the marine ecoregions of Brazil. *Front Ecol Evol* 10:2022. <https://doi.org/10.3389/fevo.2022.788879>
- Johnson KR, Owens IFP, The Global Collection Group (2023) A global approach for natural history museum collections. *Science* 379:1192–1194. <https://doi.org/10.1126/science.adf6434>
- Judkins H, Vecchione M (2013) Chapter VIII. *Doryteuthis plei*, slender inshore squid. In: Rosa R, O'Dor R, Pierce GJ (eds) *Advances in squid biology, ecology and fisheries. Part I. Myopsid squids*. Nova Science Publishers, New York, pp 241–255
- Kelly R, Fleming A, Pecl GT, von Gönner J, Bonn A (2020) Citizen science and marine conservation: a global review. *Philos Trans R Soc Lond B Biol Sci* 375:20190461. <https://doi.org/10.1098/rstb.2019.0461>
- Kowalewski M, Nawrot R, Scarponi D, Tomašových A, Zuschin M (2023) Marine conservation palaeobiology: What does the late Quaternary fossil record tell us about modern-day extinctions and biodiversity threats? *Cambridge Prisms Extinction* 1:e24. <https://doi.org/10.1017/ext.2023.22>
- Laitano MV, Fernández-Giménez AV (2016) Are mussels always the best bioindicators? Comparative study on biochemical responses of three marine invertebrate species to chronic port pollution. *Bull Environ Contam Toxicol* 97:50–55. <https://doi.org/10.1007/s00128-016-1839-y>
- Laitano MV, Nuñez JD, Cledón M (2013) Shell alterations in the limpet *Bostrycapulus odites*: a bioindicator of harbour pollution and mine residuals. *Ecol Indic* 34:345–351. <https://doi.org/10.1016/j.ecolind.2013.05.022>
- Laitano MV, Silva Barni MF, Costa PG et al (2016) Different carbon sources affect PCB accumulation by marine bivalves. *Mar Environ Res* 113:62–69. <https://doi.org/10.1016/j.marenvres.2015.11.004>
- Landro SM, Teso V, Arrighetti F (2021) The panpulmonate limpet *Siphonaria lessonii* Blainville, 1827 as a sentinel of contamination in coastal areas of Argentina. *Ecol Indic* 121:107075. <https://doi.org/10.1016/j.ecolind.2020.107075>
- Laptikhovskiy V, Arkhipkin A, Brickle P (2010) Squid as a resource shared by fish and humans on the Falkland Islands' Shelf. *Fish Res* 106:151–155. <https://doi.org/10.1016/j.fishres.2010.05.002>
- Lavrado HP, Omena EP, Bernardino AF (2017) Macrofauna bentônica do talude continental e cânions da Bacia de Campos. In: Falcão APC, Lavrado HP (eds) *Ambiente Bentônico: caracterização ambiental regional da Bacia de Campos, Atlântico Sudoeste*. Elsevier, Rio de Janeiro, pp 259–306
- Leão ZMAN, Kikuchi RKP, Testa V (2003) Corals and coral reefs of Brazil. In: Cortés J (ed) *Latin American coral reefs*. Elsevier Science, Amsterdam, pp 9–52
- Leite TS, Mather JA (2008) A new approach to octopuses' body pattern analysis: a framework for taxonomy and behavioral studies. *Am Malacol Bull* 24:31–41. <https://doi.org/10.4003/0740-2783-24.1.31>
- Leite TS, Haimovici M, Lins JE (2008a) A pesca de polvos no Arquipélago de Fernando de Noronha, Brasil. *Bol Inst Pesca* 34:271–280
- Leite TS, Haimovici M, Lins Oliveira JE (2008b) Uma proposta de manejo para a pesca do polvo *Octopus insularis* Leite & Haimovici, 2008 (Mollusca: Cephalopoda) no Arquipélago de Fernando de Noronha, Brasil. *Arq Cienc Mar* 41:81–89. <https://doi.org/10.32360/acmar.v41i1.6079>
- Leite TS, Haimovici M, Molina W, Warnke K (2008c) Morphological and genetic description of *Octopus insularis*, a new cryptic species in the *Octopus vulgaris* complex (Cephalopoda: Octopodidae) from the tropical southwestern Atlantic. *J Molluscan Stud* 74:63–74. <https://doi.org/10.1093/mollus/eym050>
- Leite TS, Haimovici M, Mather J, Oliveira JEL (2009) Habitat, distribution, and abundance of the commercial octopus (*Octopus insularis*) in a tropical oceanic island, Brazil: information for management of an artisanal fishery inside a marine protected area. *Fish Res* 98:85–91. <https://doi.org/10.1016/j.fishres.2009.04.001>
- Leite TS, Batista A, Lima F, Barbosa J, Mather J (2016) Geographic variability of *Octopus insularis* diet: from oceanic island to continental populations. *Aquat Biol* 25:17–27. <https://doi.org/10.3354/ab00655>
- Leite TS, Vidal EAG, Lima FD et al (2021) A new species of pygmy *Paroctopus* Naef, 1923 (Cephalopoda: Octopodidae): the smallest southwestern Atlantic octopod, found in sea debris. *Mar Biodivers* 51:68. <https://doi.org/10.1007/s12526-021-01201-z>
- Leite TS, Batista B, Melo CC, Vidal ÉAG, Sales JBL, Specht LM, Côrtes MO, Schroeder R, Schwarz R, Martins RS, Santos RA, Scalco ACS, Schneider F, Dolphine PM, Haimovici M (2024) *Octopus insularis*. Sistema de Avaliação do Risco de Extinção da Biodiversidade—SALVE. <https://doi.org/10.37002/salve.ficha.9943.2>
- Lima LLC, Passos FD (2021) Marine bioinvasions: bivalve molluscs introduced in northeast Brazil. *Diversitas J* 6:507–526. <https://doi.org/10.17648/diversitas-journal-v6i1-1684>
- Lima FD, Leite TS, Haimovici M, Oliveira JEL (2014a) Gonadal development and reproductive strategies of the tropical octopus (*Octopus insularis*) in northeast Brazil. *Hydrobiologia* 725(1):7–21. <https://doi.org/10.1007/s10750-013-1718-z>
- Lima FD, Leite TS, Haimovici M, Nóbrega MF, Oliveira JEL (2014b) Population structure and reproductive dynamics of *Octopus insularis* (Cephalopoda: Octopodidae) in a coastal reef environment along northeastern Brazil. *Fish Res* 152:86–92. <https://doi.org/10.1016/j.fishres.2013.08.009>
- Lima FD, Berbel-Filho WM, Leite TS, Rosas C, Lima SMQ (2017) Occurrence of *Octopus insularis* Leite and Haimovici, 2008 in the tropical northwestern Atlantic and implications of species misidentification to octopus fisheries management. *Mar Biodivers* 47:723–734. <https://doi.org/10.1007/s12526-017-0638-y>
- Lima F, Ángeles-González L, Leite TS, Lima SMQ (2020) Global climate changes over time shape the environmental niche distribution of *Octopus insularis* in the Atlantic Ocean. *Mar Ecol Prog Ser* 652:111–121. <https://doi.org/10.3354/meps13486>
- Lima-Verde FB, Castro IB, Rocha-Barreira CA (2010) Imposex occurrence in *Nassarius vibex* from South America: a potential bioindicator in estuarine environments. *Mar Biodivers Rec* 3:e30. <https://doi.org/10.1017/S1755267209990716>
- Limongi P, Ortega L, Horta S, Burone L, Carranza A (2023) Bioregionalization in a data-poor situation: mapping of Uruguayan marine benthic regions. *Front Mar Sci*. <https://doi.org/10.3389/fmars.2023.1130827>
- López MS, Coutinho R, Ferreira CEL, Rilov G (2010) Predator–prey interactions in a bioinvasion scenario: differential predation by native predators on two exotic rocky intertidal bivalves. *Mar Ecol Prog Ser* 403:101–112. <https://doi.org/10.3354/meps08409>
- Machado FM, Miranda MS, Salvador RB et al (2023) How many species of Mollusca are there in Brazil? A collective taxonomic effort to reveal this still unknown diversity. *Zoologia* 40:e23026. <https://doi.org/10.1590/S1984-4689.v40.e23026>
- Maciel DC, Castro IB, Souza JRB et al (2018) Assessment of organotins and imposex in two estuaries of the northeastern Brazilian coast. *Mar Pollut Bull* 126:473–478. <https://doi.org/10.1016/j.marpolbul.2017.11.061>
- Maggioni R, Rocha RS, Viana JT et al (2023) Genetic diversity patterns of lionfish in the southwestern Atlantic Ocean reveal a

- rapidly expanding stepping-stone bioinvasion process. *Sci Rep* 13:13469. <https://doi.org/10.1038/s41598-023-40407-y>
- Magris RA, Costa MDP, Ferreira CEL et al (2021) A blueprint for securing Brazil's marine biodiversity and supporting the achievement of global conservation goals. *Divers Distrib* 27:198–215. <https://doi.org/10.1111/ddi.13183>
- Maia RC, Coutinho R (2013) The influence of mangrove structure on the spatial distribution of *Melampus coffeus* (Gastropoda: Ellobiidae) in Brazilian estuaries. *Pan-Am J Aquat Sci* 8:21–29
- Maia RC, Tanaka MO (2007) Avaliação de efeitos de espécies de mangue na distribuição de *Melampus coffeus* (Gastropoda, Ellobiidae) no Ceará, nordeste do Brasil. *Iheringia Ser Zool* 97:379–382. <https://doi.org/10.1590/S0073-47212007000400004>
- Maia RC, Troncoso JS (2022) Evaluation of the synergistic effects of climate change on estuarine ecosystems at temperate and tropical latitudes using littorinids (Mollusca: Gastropoda) as indicators. *Braz J Anim Environ Res* 5:1642–1660. <https://doi.org/10.34188/bjaerv5n2-017>
- Marcomini SC, López RA (2006) Evolution of a beach nourishment project at Mar del Plata. *J Coast Res* 39:834–837
- Marín YH, Horta S, Chocca JF, Defeo O (2020) Historical expansion and diversification of Uruguayan fisheries in the Río de la Plata and the Atlantic Ocean: the concept of “métier” and the identification of high-intensity fishing areas. *Ocean Coast Manag* 184:104919. <https://doi.org/10.1016/j.ocecoaman.2019.104919>
- Marques RC, Breves A (2014) First record of *Pinctada imbricata* Röding, 1798 (Bivalvia: Pterioidea) attached to a rafting item: a potentially invasive species on the Uruguayan coast. *Mar Biodivers* 45:333–337. <https://doi.org/10.1007/s12526-014-0258-810.1>
- Márquez F, González-José R, Bigatti G (2011) Combined methods to detect pollution effects on shell shape and structure in neogastropods. *Ecol Indic* 11(2):248–254. <https://doi.org/10.1016/j.ecoli.2010.05.001>
- Márquez F, Primost MA, Bigatti G (2017) Shell shape as a biomarker of marine pollution historic increase. *Mar Pollut Bull* 114(2):816–820. <https://doi.org/10.1016/j.marpolbul.2016.11.018>
- Márquez F, Zabala S, Bökenhans V et al (2024) Predation of the invasive green crab *Carcinus maenas* on the edible snail *Buccinastrium deforme*, targeted as the most important nearshore marine gastropod fishery from Patagonia, Argentina. *Reg Stud Mar Sci* 69:103299. <https://doi.org/10.1016/j.risma.2023.103299>
- Martinez AS (2012) Spatial distribution of the invasive bivalve *Isognomon bicolor* on rocky shores of Arvoredo Island (Santa Catarina, Brazil). *J Mar Biol Assoc U K* 92:495–503. <https://doi.org/10.1017/S0025315411000865>
- Martinez AS, Underwood T, Christofolletti RA et al (2022) Reviewing the effects of contamination on the biota of Brazilian coastal ecosystems: Scientific challenges for a developing country in a changing world. *Sci Total Environ* 803:150097. <https://doi.org/10.1016/j.scitotenv.2021.150097>
- Martins RS, Juanicó M (2018) Biology, distribution and geographic variation of loliginid squids (Mollusca: Cephalopoda) off southwestern Atlantic. *Zoologia* 35:1–16. <https://doi.org/10.3897/zooologia.35.e23176>
- Martins VS, Schiavetti A, Souto FJB (2011) Ethnoecological knowledge of the artisan fishermen of octopi (*Octopus* spp.) in the community of Coroa Vermelha (Santa Cruz Cabrália, Bahia). *An Acad Bras Cienc* 83(2):513–522. <https://doi.org/10.1590/S0001-37652011000200011>
- Massello A, Defeo O (2016) The deep-sea red crab *Chaceon notialis* (Geryoniidae) in the southwestern Atlantic Ocean: spatial patterns and long-term effects of fishing. *Fish Res* 183:254–262. <https://doi.org/10.1016/j.fishres.2016.06.016>
- Massara Paletto V, Commendatore MG, Esteves JL (2008) Hydrocarbon levels in sediments and bivalve mollusks from Bahía Nueva (Patagonia, Argentina): an assessment of probable origin and bioaccumulation factors. *Mar Pollut Bull* 56:2100–2105. <https://doi.org/10.1016/j.marpolbul.2008.08.026>
- Mateucci SD (2012) Ecorregión Estepa Patagónica. In: Morello J, Mateucci SD, Rodríguez AF, Silva M (eds) Ecorregiones y complejos ecosistémicos argentinos. Orientación Gráfica Editora, pp 549–654
- Matthews-Cascon H, Bezerra LA, Barroso CX et al (2018) Marine benthic communities affected by the Doce River (southwestern Atlantic): baseline before a mining disaster. *Mar Pollut Bull* 135:1000–1006. <https://doi.org/10.1016/j.marpolbul.2018.08.020>
- McCauley DJ, Pinsky ML, Palumbi SR et al (2015) Marine defaunation: animal loss in the global ocean. *Science* 347:1255641. <https://doi.org/10.1126/science.1255641>
- McLachlan A, Defeo O (2018) The ecology of sandy shores. Elsevier, London
- McLachlan A, Defeo O (2023) Where will the children play? A personal perspective on sandy beaches. *Estuar Coast Shelf Sci* 280:108186. <https://doi.org/10.1016/j.ecss.2022.108186>
- Mega ER (2019) The battle to rebuild centuries of science after an epic inferno. *Nature* 571:312–315. <https://doi.org/10.1038/d41586-019-02141-2>
- Meirelles CAO, Castro ÍB, Pinheiro JCL (2008) A first record of biphallaia in imposed female of *Leucozonia nassa* (Caenogastropoda: Fasciolaridae). *Mar Biodivers Rec* 1:e39. <https://doi.org/10.1017/S1755267207004241>
- Mendez MM, Schwindt E, Bortolus A et al (2015) Ecological impacts of the austral-most population of *Crassostrea gigas* in South America: a matter of time? *Ecol Res* 30:979–987. <https://doi.org/10.1007/s11284-015-1298-7>
- Mendez MM, Livore JP, Márquez F, Bigatti G (2021) Mass mortality of foundation species on rocky shores: testing a methodology for a continental monitoring program. *Front Mar Sci*. <https://doi.org/10.3389/fmars.2021.620866>
- Messano L, Gonçalves J, Messano H, Campos S, Coutinho R (2019) First report of the Asian green mussel *Perna viridis* (Linnaeus, 1758) in Rio de Janeiro, Brazil: a new record for the southern Atlantic Ocean. *BioInvasions Rec* 8:653–660. <https://doi.org/10.3391/bir.2019.8.3.22>
- Ministério da Pesca e Aquicultura (MPA) (2011) Boletim estatístico da pesca e aquicultura. Brazil
- Mohriak W (2020) Genesis and evolution of the South Atlantic volcanic islands offshore Brazil. *Geo-Mar Lett* 40:1–33. <https://doi.org/10.1007/s00367-019-00631-w>
- Morsan EM (2003) Spatial analysis and abundance estimation of the southernmost population of purple clam, *Amiantis purpurata* in Patagonia (Argentina). *J Mar Biol Assoc U K* 83:1073–1082. <https://doi.org/10.1017/S0025315403008294h>
- Morsan E, Zaidman P, Ocampo Reinaldo M, Ciocco N (2010) Population structure, distribution and harvesting of southern geoduck, *Panopea abbreviata*, in San Matias Gulf (Patagonia, Argentina). *Sci Mar* 74:763–772. <https://doi.org/10.3989/scimar.2010.74n4763>
- Morsan EM, Sesar G, Medina Foucher C, Laco ML (2017) Assessment against MSC principles and criteria for: Patagonian scallop bottom otter trawl fishery in Argentine Sea (*Zygochlamys Patagonica*).
- Moscatelli R (1987) The superfamily Strombacea from western Atlantic. Antonio Ae Nanô & Filho, São Paulo
- Mota ELS, Alves RRN, Dias TLP (2020) Fishing, trade, and local ecological knowledge of the marine gastropod, *Cassia tuberosa*—a target species of the international shell trade. *Ethnobiol Conserv* 9:23. <https://doi.org/10.15451/ec2020-06-9.23-1-11>
- Narvarte M, González R, Fernández M (2006) Comparison of *Tehuelche octopus* (*Octopus tehuelchus*) abundance between

- an open-access fishing ground and a marine protected area: evidence from a direct development species. *Fish Res* 79:112–119. <https://doi.org/10.1016/j.fishres.2006.02.013>
- Narvarte M, González R, Filippo P (2007) Artisanal mollusk fisheries in San Matías Gulf (Patagonia, Argentina): an appraisal of the factors contributing to unsustainability. *Fish Res* 87:68–76. <https://doi.org/10.1016/j.fishres.2007.06.012>
- Nascimento NG, Silva GDV, Silva GAV, Maia RC (2022) Aspects of the population dynamics of *Anomalocardia brasiliensis*: subsidies for sustainable fishing in the west coast of Ceará, Brazil. *Cienc Anim Bras* 23:e72781. <https://doi.org/10.1590/1809-6891v23e-72781E>
- Navarro G, Rozycki V, Monsalvo M (2019) Estadísticas de la pesca marina en la Argentina: evolución de los desembarques 2012/2016. Ministerio de Agricultura, Ganadería y Pesca, Buenos Aires
- Nieto Vilela RA, Cumplido M, González Giorgis Y, Gil MN, Bigatti G (2019) Reproduction and nutritional values of the edible limpet *Nacella magellanica* (Gastropoda: Patellogastropoda). *Sci Mar* 83:237. <https://doi.org/10.3989/scimar.04825.06A>
- Nishida AK, Nordi N, Alves RRRN (2004) Abordagem etnoecológica da coleta de moluscos no litoral paraibano. *Trop Oceanogr* 32:53–68. <https://doi.org/10.5914/tropocean.v32i1.5034>
- Nishida AK, Nordi N, Alves RRRN (2006) Mollusc gathering in northeast Brazil: an ethnoecological approach. *Hum Ecol* 34:133–145. <https://doi.org/10.1007/s10745-005-9005-x>
- Norman MD, Huchberg FG (2005) The current state of octopus taxonomy. *Phuket Mar Biol Cent Res Bull* 66:127–154
- Nunes BZ, Huang Y, Ribeiro VV et al (2023a) Microplastic contamination in seawater across global marine protected areas boundaries. *Environ Pollut* 316:120692. <https://doi.org/10.1016/j.envpol.2022.120692>
- Nunes BZ, Moreira LB, Xu EG, Castro IB (2023b) A global snapshot of microplastic contamination in sediments and biota of marine protected areas. *Sci Total Environ* 865:161293. <https://doi.org/10.1016/j.scitotenv.2022.161293>
- O'Brien CE, Bennice CO, Leite TS (2021) A field guide to distinguishing *Octopus insularis* and *Octopus americanus* (Octopoda: Octopodidae). *Zootaxa* 5060:589–594. <https://doi.org/10.11646/zootaxa.5060.4.8>
- Obura DO, DeClerck F, Verburg PH et al (2023) Achieving a nature- and people-positive future. *One Earth* 6:105–117. <https://doi.org/10.1016/j.oneear.2022.11.013>
- Ojeda M, Cossi PF, Rimondino GN et al (2021) Microplastics pollution in the intertidal limpet, *Nacella magellanica*, from Beagle Channel (Argentina). *Sci Total Environ* 795:148866. <https://doi.org/10.1016/j.scitotenv.2021.148866>
- Oliva AL, Ovaert J, Arias AH, Souissi S, Marcovecchio JE (2015) Mussels as bioindicators of PAHs pollution within Argentinean coastal environments, South America. *Int J Environ Res* 9:1293–1304. <https://doi.org/10.22059/IJER.2015.1021>
- Oliveira CDC, Absalão RS (2009) The genera *Myonera*, *Octoporia*, and *Protocuspidaria* (Pelecypoda, Cuspidariidae) from deep waters of Campos Basin, Rio de Janeiro, Brazil with descriptions of two new species. *Am Malacol Bull* 27:141–156. <https://doi.org/10.4003/006.027.0212>
- Oliveira MEGC, Russo CAM, Lazoski C, Vianna PRFG, Solé-Cava AM (2005) Genetic variation and population structure of two species of neo-tropical mud-mussels (*Mytella* spp.). *Gen Mol Res* 4:197–202
- Oliveira U, Azevedo F, Dias A et al (2024) Beta diversity and regionalization of the western Atlantic marine biota. *J Biogeogr*. <https://doi.org/10.1111/jbi.14837>
- Olivier SR, Penchaszadeh PE (1968) Efectivos de almeja amarilla *Mesodesma mactroides* en las costas de la provincia de Buenos Aires y pautas para su explotación racional. Proyecto de desarrollo pesquero, serie de informes técnicos 8
- Orensanz JM, Schwindt E, Pastorino G et al (2002) No longer the pristine confines of the world ocean: a survey of exotic marine species in the southwestern Atlantic. *Biol Invasions* 4:115–143. <https://doi.org/10.1023/A:1020596916153>
- Orensanz JM, Parma AM, Jerez G et al (2005) What are the key elements for the sustainability of “S-fisheries”? Insights from South America. *Bull Mar Sci* 76:527–556
- Orensanz JM, Parma AM, Turk T, Valero J (2006) Dynamics, assessment and management of exploited natural populations, Chap 14. In: Shumway S, Parsons GJ (eds) *Scallops: biology, ecology, and aquaculture*. Elsevier Science Publishers, Amsterdam, pp 765–868
- Orensanz JM, Parma AM, Ciocco NF, Cinti A (2007) Achievements and setbacks in the commercial diving fishery of San José Gulf, Argentina Patagonia. In: McClanahan T, Castilla JC (eds) *Fisheries management: progress towards Sustainability*. Blackwell Publishing, New York, pp 68–88
- Orensanz JM, Parma AM, Ciocco NF, Cinti A (2003) Programa de entrada limitada para la pesca comercial de mariscos mediante buceo en el Golfo San José. Noveno Informe de la Comisión Técnica para la evaluación y el seguimiento de las pesquerías de Mariscos por Buceo de Chubut
- Orensanz JM, Cinti A, Parma AM, Burotto L, Espinosa Guerrero S, Sosa Cordero E, Sepúlveda C, Toral Granda V (2013) Latin American rights-based fisheries targeting sedentary resources. In: Orensanz JM, Seijo JC (eds) *Rights-based management in Latin American fisheries*. FAO Fish Aquac Tech Pap no. 582. FAO, Rome, pp 2–69
- Ortega L, Celentano E, Delgado E, Defeo O (2016) Climate change influences on abundance, individual size and body abnormalities in a sandy beach clam. *Mar Ecol Prog Ser* 545:203–213. <https://doi.org/10.3354/meps11643>
- Ortega ASB, Maranhão LA, Nobre CR et al (2019) Detoxification, oxidative stress, and cytogenotoxicity of crack cocaine in the brown mussel *Perna perna*. *Environ Sci Pollut Res* 26(27):27569–27578. <https://doi.org/10.1007/s11356-018-1600-7>
- Ortiz N, Ré ME (2011) The eggs and hatchlings of the octopus *Robsonella fontaniana* (Cephalopoda: Octopodidae). *J Mar Biol Assoc U K* 91:705–713. <https://doi.org/10.1017/S0025315410001232>
- Ortiz N, Ré ME (2019) Intertidal fishery of the Patagonian red octopus *Enteroctopus megalocyathus* (Gould, 1852): reproductive status and catch composition in the north of San Jorge Gulf (Patagonian Atlantic Coast). *J Shellfish Res* 38:619–627. <https://doi.org/10.2983/035.038.0313>
- Ortiz N, Ré ME, Márquez F, Glembocki NG (2011) The reproductive cycle of the red octopus *Enteroctopus megalocyathus* in fishing areas of northern Patagonian coast. *Fish Res* 110:217–223. <https://doi.org/10.1016/j.fishres.2011.03.016>
- Ortiz N (2009) Biología poblacional del pulpo colorado *Enteroctopus megalocyathus* en la costa patagónica norte y central y sus implicancias en el manejo pesquero. Dissertation, Universidad de Buenos Aires
- Pacheco LI, Teso V, Pastorino G (2022) Taxonomy and biogeography of bivalves of the genus *Cuspidaria* Nardo, 1840, from the southern southwestern Atlantic deep sea. *Malacologia* 65:137–175. <https://doi.org/10.4002/040.065.0109>
- Pacheco L, Teso V, Pastorino G (2024) Use of traditional tools and micro-computed tomography for the taxonomy of carnivorous bivalves from the deep waters of southwestern Atlantic. *Zool J Linnean Soc* 202(1):zlae118. <https://doi.org/10.1093/zoolinnean/zlae118>
- Padula V, Araujo AK, Matthews-Cascon H, Schrödl M (2014) Is the Mediterranean nudibranch *Cratena peregrina* (Gmelin, 1791)

- present on the Brazilian coast? Integrative species delimitation and description of *Cratena minor* n. sp. J Molluscan Stud 80:575–584. <https://doi.org/10.1093/mollus/eyu052>
- Parker G, Paterlini CM, Violante RA, Costa IP, Marcolini SI, Cavallotto JL (1999) Descripción geológica de la terraza rioplatense (plataforma interior del noreste Bonaerense). Servicio Geológico y Minero Argentino, Buenos Aires, boletín no. 273
- Passos FD, Miranda MS, Corrêa PVF (2019) Synopsis of the knowledge on the Brazilian aplacophorans (Mollusca: Caudofoveata & Solenogastres). Biota Neotrop 19:e20180545. <https://doi.org/10.1590/1676-0611-BN-2018-0545>
- Passos FD, Corrêa PVF, Miranda MS (2021) Better alone than in bad company: new species of caudofoveate limifossorid (Mollusca, Aplacophora) unravels patterns of distribution hidden in the deep Atlantic. Front Mar Sci. <https://doi.org/10.3389/fmars.2021.669478>
- Passos FD, Fassina PV, Corrêa PVF, Miranda MS (2022) Separated from the cradle: a new species of *Falcidens* (Mollusca, Aplacophora, Caudofoveata) reveals weird patterns of distribution in the deep-sea. Deep-Sea Res I 186:103825. <https://doi.org/10.1016/j.dsr.2022.103825>
- Pastorino G (2016a) Revision of the genera *Pareuthria* Strebel, 1905, *Glypteuthria* Strebel, 1905 and *Meteuthria* Thiele, 1912 (Gastropoda: Buccinulidae) with the description of three new genera and two new species from southwestern Atlantic waters. Zootaxa 4179:301–344. <https://doi.org/10.11646/zootaxa.4179.3.1>
- Pastorino G (2016b) First report of the family Laubierinidae Warén & Bouchet, 1990 (Gastropoda: Tonnoidea) in the southwestern Atlantic. Molluscan Res 36:108–111. <https://doi.org/10.1080/13235818.2015.1128582>
- Pastorino G (2019) A new deep water gastropod of the genus *Parabuccinum* (Neogastropoda: Buccinulidae) from southwestern Atlantic waters with new data on the distribution of all species. Mar Biodivers 49:913–922. <https://doi.org/10.1007/s12526-018-0876-7>
- Pastorino G, Chiesa IL (2014) The family Caecidae (Gastropoda: Caenogastropoda) in Argentine waters. Nautilus 128:40–50
- Pastorino G, Sánchez N (2016) Southwestern Atlantic species of conoidean gastropods of the genus *Aforia* Dall, 1889. Zootaxa 4109:458–470. <https://doi.org/10.11646/zootaxa.4109.4.4>
- Pastorino G, Scarabino F (2008) Two new deep-sea muricids (Gastropoda) from Argentina. Nautilus 122:107–114
- Pastorino G, Penchaszadeh PE, Schejter L, Bremec C (2000) *Rapana venosa* (Valenciennes, 1846) (Mollusca: Muricidae): a new gastropod in south Atlantic waters. J Shellfish Res 19:897–899
- Pazos RS, Spaccesi F, Gómez N (2020) First record of microplastics in the mussel *Limnoperna fortunei*. Reg Stud Mar Sci 38:101360. <https://doi.org/10.1016/j.rsma.2020.101360>
- Peake J, Bogdanoff AK, Layman CA et al (2018) Feeding ecology of invasive lionfish (*Pterois volitans* and *Pterois miles*) in the temperate and tropical western Atlantic. Biol Invasions 20:2567–2597. <https://doi.org/10.1007/s10530-018-1720-5>
- Pecl GT, Araújo MB, Bell JD et al (2017) Biodiversity redistribution under climate change: impacts on ecosystems and human well-being. Science 355:eaai9214. <https://doi.org/10.1126/science.aai9214>
- Pedro NC, Salvador RB, Simone LRL (2023) First record of the exotic *Indothais lacera* (Gastropoda, Muricidae) in Brazil. Pap Avulsos Zool 63:e202363004. <https://doi.org/10.11606/1807-0205/2023.63.004>
- Penchaszadeh PE, Averbuj A, Cledón M (2001) Imposex in gastropods from Argentina (south-western Atlantic). Mar Pollut Bull 42:790–791. [https://doi.org/10.1016/S0025-326X\(01\)00098-4](https://doi.org/10.1016/S0025-326X(01)00098-4)
- Penchaszadeh PE, Atencio M, Martinez MI, Pastorino G (2016) Giant egg capsules and hatchlings in a deep-sea moon snail (Naticidae) from a southwestern Atlantic canyon. Mar Biol 163:209. <https://doi.org/10.1007/s00227-016-2990-z>
- Penchaszadeh PE, Teso V, Pastorino G (2017) Spawn in two deep-sea volute gastropods (Neogastropoda: Volutidae) from southwestern Atlantic waters. Deep-Sea Res I 130:55–62. <https://doi.org/10.1016/j.dsr.2017.10.011>
- Penchaszadeh PE, Pastorino G, Martinez MI, Miloslavich P (2019) Spawn and development of the gastropod *Americominella longisetosa* (Castellanos and Fernández, 1972) (Mollusca: Buccinidae) from the southwestern Atlantic deep sea. Deep-Sea Res I 143:43–49. <https://doi.org/10.1016/j.dsr.2018.11.011>
- Penchaszadeh P, Arrighetti F, Aldea C, Teso V (2022) Reproductive pattern of *Trochita pileolus* (d'Orbigny, 1841) (Gastropoda: Calyptraeidae) and different type of intracapsular development in *Trochita* species from South America. Polar Biol 45:59–69. <https://doi.org/10.1007/s00300-021-02972-8>
- Pereira OM, Machado IC, Henriques MB, Galvão MSN, Yamanaka N (2001) Stock evaluation of the oyster *Crassostrea brasiliana* in mangroves of rivers and tidal creeks ('gamboas') in Cananéia lagoon estuarine system (São Paulo State, Brazil). Bol Inst Pesca 27:85–95
- Pereira OM, Hilberath RC, Ansarah PRAC, Galvão MSN (2003) Estimativa da produção de *Mytella falcata* e de *M. guyanensis* em bancos naturais do estuário de Ilha Comprida—SP—Brasil. Bol Inst Pesca 29:139–149
- Perez JAA, Haimovici M (1991) Sexual maturation and reproductive cycle of *Eledone massyae* Voss 1964 (Cephalopoda: Octopodidae) in southern Brazil. Bull Mar Sci 49:270–279
- Perez JAA, de Aguiar DC, Oliveira UC (2002) Biology and population dynamics of the long-finned squid *Loligo plei* (Cephalopoda: Loliginidae) in southern Brazilian waters. Fish Res 58(3):267–279. [https://doi.org/10.1016/S0165-7836\(01\)00397-6](https://doi.org/10.1016/S0165-7836(01)00397-6)
- Perez JAA, Martins RS, Santos RA (2004) Cefalópodes capturados pela pesca comercial de talude no Sudeste e Sul do Brasil. Notas Tec FACIMAR 8:65–74
- Perez JAA, Silva TN, Schroeder R, Schwarz R, Martins RS (2009) Biological patterns of the Argentine shortfin squid *Illex argentinus* in the slope trawl fishery off Brazil. Lat Am J Aquat Res 37(3):409–428
- Perez JAA, Alves ES, Clark MR et al (2012) Patterns of life on the southern Mid-Atlantic Ridge: compiling what is known and addressing future research. Oceanogr 25:16–31. <https://doi.org/10.5670/oceanogr.2012.102>
- Pérez AF, Ojeda M, Rimondino GN et al (2020) First report of microplastics presence in the mussel *Mytilus chilensis* from Ushuaia Bay (Beagle Channel, Tierra del Fuego, Argentina). Mar Pollut Bull 161:111753. <https://doi.org/10.1016/j.marpolbul.2020.111753>
- Petracco M, Camargo RM, Berenguel TA et al (2015) Evaluation of the use of *Olivella minuta* (Gastropoda, Olividae) and *Hastula cinerea* (Gastropoda, Terebridae) as TBT sentinels for sandy coastal habitats. Environ Monit Assess 187:440. <https://doi.org/10.1007/s10661-015-4650-z>
- Pezzuto PR, Borzone CA (2004) The collapse of the scallop *Euvola ziczac* (Linnaeus, 1758) (Bivalvia: Pectinidae) fishery in Brazil: changes in distribution and relative abundance after 23 years of exploitation. Braz J Oceanogr 52:225–236
- Pezzuto PR, Schio C, Almeida TC (2010) Efficiency and selectivity of the *Anomalocardia brasiliana* (Mollusca: Veneridae) hand dredge used in southern Brazil. J Mar Biol Assoc U K 90(7):1455–1464. <https://doi.org/10.1017/S0025315410000317>
- Pimenta AD, Andrade BG, Absalão RS (2018) Taxonomy of Nystiellidae (Caenogastropoda, Epitonioidea) from Brazil, with description of three new species from the south-western Atlantic. J Mar Biol Assoc U K 98:1619–1644. <https://doi.org/10.1017/S0025315417000984>

- Poloczanska ES, Brown CJ, Sydeman WJ et al (2013) Global imprint of climate change on marine life. *Nat Clim Change* 3:919–925. <https://doi.org/10.1038/nclimate1958>
- Póvoa AA, Skinner LF, Araújo FV (2021) Fouling organisms in marine litter (rafting on abiogenic substrates): a global review of literature. *Mar Pollut Bull* 166:112189. <https://doi.org/10.1016/j.marpolbul.2021.112189>
- Praxedes AGP, Castro DL, Torres LC, Gambôa LAP, Hackspacher PC (2019) New insights of the tectonic and sedimentary evolution of the Rio Grande Rise, South Atlantic Ocean. *Mar Pet Geol* 10:335–346. <https://doi.org/10.1016/j.marpetgeo.2019.07.035>
- Primost MA, Averbuj A, Bigatti G (2015) Variability of imposex development and reproductive alterations in the Patagonian gastropod *Buccinanops globulosus* inhabiting a polluted harbour area. *Rev Mus Argent Cienc* 17:167–171
- Primost MA, Sabatini SE, Di Salvatore P, Ríos De Molina MC, Bigatti G (2017) Oxidative stress indicators in populations of the gastropod *Buccinanops globulosus* affected by imposex. *J Mar Biol Assoc U K* 97:35–42. <https://doi.org/10.1017/S0025315415002155>
- Primost MA, Commendatore M, Torres PJ, Bigatti G (2018) PAHs contamination in edible gastropods from north Patagonian harbor areas. *Mar Pollut Bull* 135:828–831. <https://doi.org/10.1016/j.marpolbul.2018.08.021>
- Queiroz LR, Castro IB, Rocha-Barreira CA (2007) New imposex development index (IDI) for *Stramonita haemastoma* (Mollusca: Muricidae): a transplantation experiment in the Brazilian north-east. *J Braz Soc Ecotox* 2:249–256
- Rabone M, Wiethase JH, Simon-Lledó E et al (2023) How many metazoan species live in the world's largest mineral exploration region? *Curr Biol* 33(12):2383–2396.e5. <https://doi.org/10.1016/j.cub.2023.04.052>
- Ré ME (1998a) *Pulpos octopódidos* (Cephalopoda, Octopodidae). In: Boschi E (ed) *El Mar Argentino y sus recursos pesqueros*, Tomo 2. Publicaciones especiales INIDEP, Mar del Plata, pp 69–98
- Ré ME (1998b) *Pesquerías de pulpos*. In: Boschi E (ed) *El Mar Argentino y sus recursos pesqueros*, Tomo 2. Publicaciones especiales INIDEP, Mar del Plata, pp 99–114
- Ré ME, Taylor R (1981) La pesca de pulpos en Argentina. *Artes de captura utilizadas y estadísticas pesqueras hasta 1978*. Centro Nacional Patagónico-CONICET, N° 52
- Ribeiro VV, Nobre CR, Moreno BB et al (2023) Oysters and mussels as equivalent sentinels of microplastics and natural particles in coastal environments. *Sci Total Environ* 874:162468. <https://doi.org/10.1016/j.scitotenv.2023.162468>
- Ribeiro-Brasil DRG, Castro IB, Petracco M et al (2021) Spatial distribution of butyltins and imposex in eastern Brazilian Amazon. *Mar Pollut Bull* 165:112155. <https://doi.org/10.1016/j.marpolbul.2021.112155>
- Riestra G, Fabiano G, Santana O (2000) El caracol negro (*Adelomelon brasiliense*) como recurso no tradicional de importancia para el país: análisis socio-económico de la pesquería y medidas precautorias de manejo. In: Rey M (ed) *Recursos pesqueros no tradicionales: moluscos bentónicos marinos*, Uruguay, pp 82–92
- Ríos MF, Hernández-Moresino RD, Galván DE (2020) Assessing urban microplastic pollution in a benthic habitat of Patagonia Argentina. *Mar Pollut Bull* 159:111491. <https://doi.org/10.1016/j.marpolbul.2020.111491>
- Rocha LM (2013) *Ecología humana e manejo participativo da pesca do búzio Anomalocardia brasiliense* (Gmelin, 1791) (Bivalvia: Veneridae) na Reserva de Desenvolvimento Sustentável Estadual Ponta do Tubarão (RN). Universidade Federal do Rio Grande do Norte, Natal
- Rockström J, Steffen W, Noone K et al (2009) A safe operating space for humanity. *Nature* 461:472–475. <https://doi.org/10.1038/461472a>
- Rodrigues JAM, Santos MR, Maia RC (2020) Estudo do imposex no gastrópode *Stramonita brasiliensis* (Claremont & Reid, 2011) nos municípios de Acaraú e Itarema, litoral oeste do Ceará, Brasil. *Arq Cienc Mar* 53:126–133. <https://doi.org/10.32360/acmar.v53i1.43641>
- Rosa R, Trübenbach K, Repolho T et al (2013) Lower hypoxia thresholds of cuttlefish early life stages living in a warm acidified ocean. *Proc R Soc B Biol Sci* 280:20131695. <https://doi.org/10.1098/rspb.2013.1695>
- Rosas-Luis R, Sánchez P, Portela JM, del Rio JL (2014) Feeding habits and trophic interactions of *Doryteuthis gahi*, *Illex argentinus* and *Onykia ingens* in the marine ecosystem off the Patagonian Shelf. *Fish Res* 152:37–44. <https://doi.org/10.1016/j.fishres.2013.11.004>
- Rossato M, Castro IB, Pinho GLL (2014) Imposex in *Stramonita haemastoma*: a preliminary comparison between waterborne and dietborne exposure. *Ecotoxicol Environ Contam* 9:87–92. <https://doi.org/10.5132/eec.2014.01.011>
- Rossato M, Castro IB, Paganini CL et al (2016) Sex steroid imbalances in the muricid *Stramonita haemastoma* from TBT contaminated sites. *Environ Sci Pollut Res* 23:7861–7868. <https://doi.org/10.1007/s11356-015-5942-0>
- Ruiz MD, Iriel A, Yusseppone MS et al (2018) Trace metals and oxidative status in soft tissues of caged mussels (*Aulacomya atra*) on the north Patagonian coastline. *Ecotoxicol Environ Saf* 155:152–161. <https://doi.org/10.1016/j.ecoenv.2018.02.064>
- Sales JBL, Haimovici M, Ready JS et al (2019) Surveying cephalopod diversity of the Amazon reef system using samples from red snapper stomachs and description of a new genus and species of *octopus*. *Sci Rep* 9:5956. <https://doi.org/10.1038/s41598-019-42464-8>
- Sales JBL, Leite T, Melo CC et al (2024) *Lepidoctopus joaquina* Haimovici & Sales, 2019. Sistema de Avaliação do Risco de Extinção da Biodiversidade—SALVE. Available in: <https://salve.icmbio.gov.br>. Accessed on 19 mar 2024. <https://doi.org/10.37002/salve.ficha.38988.1>
- Sánchez N, Pastorino G (2020) The North Atlantic conoidean gastropod *Theta lyronuclea* (Raphitomidae) in deep-waters of the southwestern Atlantic. *Malacologia* 63:33–40. <https://doi.org/10.4002/040.063.0104>
- Sánchez N, Pastorino G, Penchaszadeh PE (2018) Giant eggs in the gastropod *Aforia obesa* (Conoidea: Cochlespiridae) in southwestern Atlantic deep-waters. *Zool Anz* 276:94–99. <https://doi.org/10.1016/j.jcz.2018.07.006>
- Sánchez N, Damborenea C, Pastorino G (2023) Unravelling the conoidean gastropods assigned to the genus *Propebela* (Gastropoda: Mangeliidae) from south-western Atlantic deep waters. *J Nat Hist* 57:243–256. <https://doi.org/10.1080/00222933.2023.2174056>
- Sánchez NC, Vogler RE, Beltramino AA, Pastorino G (2024) How many species of the gastropod genus *Leucosyrinx* (Conoidea: Pseudomelatomidae) are truly living in southwestern Atlantic deep-waters? *Zool Anz* 313:37–48. <https://doi.org/10.1016/j.jcz.2024.08.004>
- Sánchez N, Aldea C, Pastorino G (2025) The conoidean genus *Belalora* (Gastropoda: Mangeliidae) under the magnifying glass: an exclusive inhabitant of Magellanic waters. *Zoosystema* 47:125–132. <https://doi.org/10.5252/zoosystema2025v47a8>
- Sánchez RP, Navarro G, Rozycki V (2012) *Estadísticas de la pesca marina en la Argentina: evolución de los desembarques 1898–2010*. Ministerio de Agricultura, Ganadería y Pesca, Buenos Aires
- Sánchez-Carnero N, Góngora ME, Álvarez M, Parma A (2022) La pesca artesanal en Argentina: caminando las costas del país
- Santana MF, Ascer LG, Custódio MR, Moreira FT, Turra A (2016) Microplastic contamination in natural mussel beds from a Brazilian urbanized coastal region: rapid evaluation through

- bioassessment. *Mar Pollut Bull* 106:183–189. <https://doi.org/10.1016/j.marpolbul.2016.02.074>
- Santos CMD, Carbayo F (2021) Taxonomy as a political statement: the Brazilian case. *Zootaxa* 5047:92–94. <https://doi.org/10.11646/zootaxa.5047.1.8>
- Santos MR, Maia RC (2021) Obras de urbanização e os impactos ambientais sobre a macrofauna bentônica: estudo de caso em uma praia do litoral cearense. *J Integr Coast Zone Manag* 21:135–145. <https://doi.org/10.5894/rgci-n348>
- Santos MR, Maia RC (2023) Imposex in *Stramonita brasiliensis* (Mollusca: Gastropoda) in port terminal of Ceará: environmental assessment and perception. *Rev Aedis Ingen Cienc Ambien Invest Desarrollo Práctica* 16:1013–1029. <https://doi.org/10.22201/iingen.0718378xe.16.3.83883>
- Santos WCR, Chagas RA, Barros MRF et al (2021) Lista preliminar de bycatch de molusco em pescaria de arrasto na plataforma continental amazônica. *Arq Cienc Mar* 54:150–162. <https://doi.org/10.32360/acmar.v54i1.42007>
- Santos FS, Neves RAF, Crapez MAC, Teixeira VL, Krepesky N (2022) How does the brown mussel *Perna perna* respond to environmental pollution? A review on pollution biomarkers. *J Environ Sci* 111:412–428. <https://doi.org/10.1016/j.jes.2021.04.006>
- Sauer WHH, Gleadall IG, Downey-Breedt N et al (2021) World octopus fisheries. *Rev Fish Sci Aquacult* 29:279–429. <https://doi.org/10.1080/23308249.2019.1680603>
- Scarabino F (2004) Conservación de la malacofauna uruguaya. *Comun Soc Malacol Urug* 8:267–273
- Scarabino F, Zelaya DG, Orensanz JM et al (2015) Cold, warm, temperate and brackish: bivalve biodiversity in a complex oceanographic scenario (Uruguay, southwestern Atlantic). *Am Malacol Bull* 33:284–301. <https://doi.org/10.4003/006.033.0219>
- Scarabino F, Laporta C, Vélez Rubio G, Lagos X, Perez L, Serra S, Verrastro N (2020) Ficha costera caracol negro y sus ovi-cápsulas. Grupo Costa+ y CEIMER, Caja Costera. <https://www.costamas.edu.uy/la-caja-costera/>
- Scarabino F (2021) Las especies también son patrimonio: un análisis integral y aplicado de los valores y usos de los moluscos de un sector de la costa rochense (Uruguay). Master's thesis, Universidad de la República, Uruguay. <https://www.colibri.udelar.edu.uy/jspui/handle/20.500.12008/37632>
- Scherner F, Horta PA, de Oliveira EC et al (2013) Coastal urbanization leads to remarkable seaweed loss and community shifts along the SW Atlantic. *Mar Pollut Bull* 76(1–2):106–115. <https://doi.org/10.1016/j.marpolbul.2013.09.019>
- Schuab JM, De Paula MS, Ocaris ERY et al (2023) First record of microplastic in the Brazilian sea hare *Aplysia brasiliensis* Rang, 1828 (Mollusca: Aplysiidae). *Sci Total Environ* 895:165156. <https://doi.org/10.1016/j.scitotenv.2023.165156>
- Schwindt E, Carlton JT, Orensanz JM, Scarabino F, Bortolus A (2020) Past and future of the marine bioinvasions along the southwestern Atlantic. *Aquat Invasions* 15:11–29. <https://doi.org/10.3391/ai.2020.15.1.02>
- Siegwald J, Pastorino G, Oskars T, Malaquias MAE (2020) A new species of the deep-sea genus *Scaphander* (Gastropoda, Cephalaspidea) from the Mar del Plata submarine canyon off Argentina. *Bull Mar Sci* 96:111–126. <https://doi.org/10.5343/bms.2019.0069>
- Signorelli JH, Crespo E (2017) First record of the genus *Adipicola* (Mollusca: Bivalvia: Mytilidae) and description of a new species from the Argentine SW Atlantic Ocean. *Zootaxa* 4318:325–338. <https://doi.org/10.11646/zootaxa.4318.2.6>
- Signorelli J, Pastorino G (2015) A new species of *Laubiericoncha* (Bivalvia: Vesicomidae) from deep waters off Argentina. *Malacologia* 58:349–360. <https://doi.org/10.4002/040.058.0213>
- Sigwart JD, Chen C, Marsh L (2017) Is mining the seabed bad for mollusks? *Nautilus* 131:43–49
- Silva JB, Pereira ECG, Torres MFA (2007) Estuário de Itapessoca, Pernambuco: relação entre pescadores e pescadeiras, e as espécies de maior importância comercial. *Bol Tec Cient CEPENE* 15:81–88
- Simone LRL, Gonçalves EP (2006) Anatomical study on *Myoforceps aristatus*, an invasive boring bivalve in S.E. Brazilian coast (Mytilidae). *Pap Avulsos Zool* 46:57–65. <https://doi.org/10.1590/S0031-10492006000600001>
- Soares MO, Rabelo EF (2023) Severe ecological impacts caused by one of the worst orphan oil spills worldwide. *Mar Environ Res* 187:105936. <https://doi.org/10.1016/j.marenvres.2023.105936>
- Soares MO, Xavier FRL, Dias NM et al (2022) Alien hotspot: Benthic marine species introduced in the Brazilian semiarid coast. *Mar Pollut Bull* 174:113250. <https://doi.org/10.1016/j.marpolbul.2021.113250>
- Socrate J, Verón E, Chaparro M (2024) Public perception of offshore hydrocarbon activities in the North Argentine Basin: a study in Buenos Aires Province, Argentina. *J Coast Conserv* 28:77. <https://doi.org/10.1007/s11852-024-01078-7>
- Soria G, Orensanz JM, Morsán EM, Parma AM, Amoroso RO (2016) Scallops biology, fisheries, and management in Argentina. *Dev Aquacult Fish Sci* 40:1019–1046. <https://doi.org/10.1016/B978-0-444-62710-0.00025-0>
- Souza LS, Araújo ICV, Caetano CHS (2013) A commented list of Scaphopoda (Mollusca) found along the Brazilian coast, with two new synonymies in the genus *Gadila* Gray, 1847. *Biota Neotrop* 13:227–235. <https://doi.org/10.1590/S1676-06032013000200022>
- Souza LS, Caetano CHS, Scarabino F, Costa PMS (2020) New records and a new species of Scaphopoda (Mollusca) from the southwestern Atlantic Ocean. *Iheringia Ser Zool* 110:e2020023. <https://doi.org/10.1590/1678-4766e2020023>
- Souza BHM, Passos FD, Shimabukuro M, Sumida PYG (2021) An integrative approach distinguishes three new species of Abyssochrysoidea (Mollusca: Caenogastropoda) associated with organic falls of the deep south-west Atlantic. *Zool J Linn Soc* 191:748–771. <https://doi.org/10.1093/zoolinnean/zlaa059>
- Spotorno-Oliveira P, Coutinho R, Tâmega FTS (2018) Recent introduction of non-indigenous vermetid species (Mollusca, Vermetidae) to the Brazilian coast. *Mar Biodivers* 48:1931–1941. <https://doi.org/10.1007/s12526-017-0702-7>
- Spotorno-Oliveira P, Lopes RP, Larroque A et al (2020) First detection of the non-indigenous gastropod *Rapana venosa* in the southern-most coast of Brazil. *Cont Shelf Res* 194:104047. <https://doi.org/10.1016/j.csr.2020.104047>
- Staichak G, Ferreira-Jr AL, Moreschi Silva AC et al (2021) Bivalves with potential for monitoring microplastics in South America. *Case Stud Chem Environ Eng* 4:100119. <https://doi.org/10.1016/j.cscee.2021.100119>
- Steffen W, Broadgate W, Deutsch L, Gaffney O, Ludwig C (2015a) The trajectory of the Anthropocene: the great acceleration. *Anthr Rev* 2:81–98. <https://doi.org/10.1177/2053019614564785>
- Steffen W, Richardson K, Rockström J et al (2015b) Planetary boundaries: guiding human development on a changing planet. *Science* 347:1259855. <https://doi.org/10.1126/science.1259855>
- Storero LP, Narvarte MA, González RA (2013) Marine protected areas: reserve effect or natural variability? The Patagonian octopus case. *J Mar Biol Assoc U K* 93:259–266. <https://doi.org/10.1017/S0025315412000525>
- Storero LP (2010) Características ecológicas del pulpito *Octopus tehuelchus* en tres ambientes del Golfo San Matías. *Dissertation, Universidad Nacional de Córdoba*
- Tanaka MO, Maia RC (2006) Shell morphological variation of *Littoraria angulifera* among and within mangroves in NE Brazil. *Hydrobiologia* 559:193–202. <https://doi.org/10.1007/s10750-005-1449-x>

- Teixeira L, Creed J (2020) A decade on: an updated assessment of the status of marine non-indigenous species in Brazil. *Aquat Invasions* 15:30–43. <https://doi.org/10.3391/ai.2020.15.1.03>
- Teso V, Penchaszadeh PE (2009) Beach filling and imposex in *Olivancillaria deshayesiana* (Mollusca: Gastropoda: Olividae) from the coast of Mar del Plata, Argentina. *J Mar Biol Assoc U K* 89:557–562. <https://doi.org/10.1017/S0025315409002896>
- Teso V, Penchaszadeh PE (2019) Development of the gastropod *Trochita pileus* (Calyptaeidae) in the sub-Antarctic southwestern Atlantic. *Polar Biol* 42:171–178. <https://doi.org/10.1007/s00300-018-2412-4>
- Teso V, Arrighetti F, Penchaszadeh PE (2012) Reproductive cycle in an imposexed population of *Olivancillaria deshayesiana* (Gastropoda: Olividae) from Mar del Plata. *Argentina Aquat Biol* 15(2):111–119. <https://doi.org/10.3354/ab00407>
- Teso V, Urteaga D, Pastorino G (2019) Assemblages of certain benthic molluscs along the southwestern Atlantic: from subtidal to deep sea. *BMC Ecol* 19:1–7. <https://doi.org/10.1186/s12898-019-0263-7>
- Teso V, Martínez M, Lauretta D et al (2020) Protected development in deep-sea invertebrates. A case of study in the southwestern Atlantic Ocean deep-sea. *Environ Coast Offshore Deep-Sea Special Edition*:48–51
- Thompson C, Bacha L, Paz CHC et al (2023) Collapse of scallop *Nodipecten nodosus* production in the tropical southeast Brazil as a possible consequence of global warming and water pollution. *Sci Total Environ* 904:166873. <https://doi.org/10.1016/j.scitotenv.2023.166873>
- Todt C (2013) Aplacophoran mollusks—still obscure and difficult? *Am Malacol Bull* 31:181–187. <https://doi.org/10.4003/006.031.0110>
- Truchet DM, López ADF, Arduzzo MG et al (2021) Microplastics in bivalves, water and sediments from a touristic sandy beach of Argentina. *Mar Pollut Bull* 173:113023. <https://doi.org/10.1016/j.marpolbul.2021.113023>
- United Nation (2018) Regional agreement on access to information, public participation and justice in environmental matters in Latin America and the Caribbean. Available at: <https://treaties.un.org/doc/Treaties/2018/03/20180312%2003-04%20PM/CTC-XXVII-18.pdf>. Access: 15 Apr 2024
- United Nations (2024a) Department of Economic and Social Affairs—Sustainable Development. Available at: <https://sdgs.un.org/goals>. Accessed: 8 Jan 2024
- United Nations (2024b) UN Ocean Conference, Lisbon, Portugal. Available at: <https://www.un.org/en/conferences/ocean2022>. Accessed: 8 Jan 2024
- Urteaga D (2011) Taxonomía de los quitones (Mollusca, Polyplacophora) del Atlántico sudoccidental. Dissertation, Universidad de Buenos Aires
- Vales DG, Hernández-Orts JS, Durante CA et al (2017) *Argonauta nodosus* (Cephalopoda: Argonautidae) in Patagonian waters. *Malacologia* 62:367–371. <https://doi.org/10.4002/040.062.0211>
- Veloso VG, Sallorenzo IA, Fernandes VB (2017) Macrofauna bentônica da plataforma continental da Bacia de Campos. In: Falcão APC, Lavrado HP (eds) *Ambiente Bentônico: caracterização ambiental regional da Bacia de Campos, Atlântico Sudoeste*. Elsevier, Rio de Janeiro, pp 227–258
- Viana MG, Lima MSP, Martínez AS et al (2022) Marine fish and benthic biota before the 2019 oil spill: a baseline dataset for monitoring programs and impact assessments at Rio Grande Norte state, northeastern Brazil. *An Acad Bras Cien* 94:e20210536. <https://doi.org/10.1590/0001-3765202120210536>
- Vidal EAG, Marian JEAR, Martins RS (2013) Chapter IX: *Doryteuthis sanpaulensis*, São Paulo Squid. In: Rosa R, O’Dor R, Pierce GJ (eds) *Advances in squid biology, ecology and fisheries. Part I. Myopsid squids*. Nova Science Publishers, New York, pp 257–300
- Vieira KS, Batista Neto JA, Crapez MAC et al (2021) Occurrence of microplastics and heavy metals accumulation in native oysters *Crassostrea gasar* in the Paranaguá estuarine system, Brazil. *Mar Pollut Bull* 166:112225. <https://doi.org/10.1016/j.marpolbul.2021.112225>
- Violante RA, Laprida C, García Chaporí NL (2017) The Argentina Continental Margin: location and significance. In: Lohmann G, Mysak LA, Notholt J, Rabassa J, Unnithan V (eds) *The Argentina Continental Margin: a potential paleoclimatic-paleoceanographic archive for the southern Ocean*. Springer Briefs in Earth System Sciences. Springer, pp 11–32
- Young AM, Elliott JA (2019) Life history and population dynamics of green crabs (*Carcinus maenas*). *Fishes* 5:4. <https://doi.org/10.3390/fishes5010004>
- Zamudio KR, Kellner A, Serejo C et al (2018) Lack of science support fails Brazil. *Science* 361(6409):1322–1323. <https://doi.org/10.1126/science.aav3296>
- Zelaya DG (2015) Marine bivalves from the Argentine Coast and Continental Shelf: species diversity and assessment of the historical knowledge. *Am Malacol Bull* 33:245. <https://doi.org/10.4003/006.033.0204>

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.

Authors and Affiliations

Cristiane Xerez Barroso¹ · Eliane Pintor de Arruda² · Florencia Arrighetti³ · Andrés Averbuj⁴ · Gregorio Bigatti^{4,5} · Ramiro Braga^{6,7} · Alvar Carranza^{8,9} · Susete Wambier Christo¹⁰ · Mariana Osório Côrtes¹¹ · Paulo Márcio Santos Costa¹² · Mariano Cumplido⁴ · Omar Defeo¹³ · Augusto Luiz Ferreira Jr.¹⁴ · Manuel Haimovici¹⁵ · Tatiana Silva Leite¹⁶ · Fabrizio Marcondes Machado¹⁷ · Rafaela Camargo Maia¹⁸ · Inês Xavier Martins¹⁹ · Helena Matthews-Cascon¹ · Marcel Sabino Miranda^{17,20,21} · Nicolás Ortiz^{6,22} · Vinicius Padula²³ · Maria Gabriela Palomo³ · Flávio Dias Passos¹⁷ · Guido Pastorino³ · Natan Carvalho Pedro¹¹ · Pablo E. Penchaszadeh³ · Fabrizio Scarabino²⁴ · Javier H. Signorelli⁴ · Leonardo Santos de Souza²⁵ · Paula Spotorno²⁶ · Lenita de Freitas Tallarico²⁷ · Diego Urteaga³ · Silvina Van der Molen⁶ · Valeria Teso³

✉ Valeria Teso
valeriateso@gmail.com

¹ Laboratório de Invertebrados Marinhos do Ceará, Departamento de Biologia, Centro de Ciências, Universidade Federal do Ceará, Fortaleza, CE, Brazil

² Laboratório de Estudos em Macroinvertebrados Bentônicos, Departamento de Biologia, Centro de Ciências Humanas e Biológicas, Universidade Federal de São Carlos, Campus Sorocaba, Sorocaba, SP, Brazil

³ Laboratorio de Ecosistemas Marinos, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” (MACNBR-CONICET), Buenos Aires, Argentina

⁴ Laboratorio de Reproducción y Biología Integrativa de Invertebrados Marinos (LARBIM), Instituto de Biología de Organismos Marinos (IBIOMAR) (CCT CONICET-CENPAT), Puerto Madryn, Chubut, Argentina

⁵ Universidad Espíritu Santo, Samborondón, Ecuador

⁶ Laboratorio de Cefalópodos, Instituto de Biología de Organismos Marinos (IBIOMAR) (CCT CONICET-CENPAT), Puerto Madryn, Chubut, Argentina

⁷ Laboratorio de Crustáceos y Ecosistemas Costeros (CADIC-CONICET), Ushuaia, Tierra del Fuego, Argentina

⁸ Departamento de Ecología y Gestión Ambiental, Centro Universitario Regional del Este (CURE), Sede Maldonado, Universidad de La República, Punta del Este, Uruguay

⁹ Museo Nacional de Historia Natural, Montevideo, Uruguay

¹⁰ Programa de Pós-Graduação em Biologia Evolutiva, Departamento de Biologia Estrutural, Molecular e Genética, Departamento de Biologia Geral, Universidade Estadual de Ponta Grossa, Ponta Grossa, PR, Brazil

¹¹ Laboratório de Malacologia, Museu de Zoologia da Universidade de São Paulo (MZUSP), São Paulo, SP, Brazil

¹² Diretoria de Pesquisa e Produção, Fundação Instituto de Pesca do Estado do Rio de Janeiro (FIPERJ), Niterói, RJ, Brazil

¹³ Facultad de Ciencias, Laboratorio de Ciencias del Mar (UNDECIMAR), Montevideo, Uruguay

¹⁴ Programa de Pós-Graduação em Genética Evolutiva e Biologia Molecular, Universidade Federal de São Carlos, São Carlos, SP, Brazil

¹⁵ Laboratório de Dinâmica de Populações Pesqueiras, Instituto de Oceanografia, Universidade Federal do Rio Grande, Rio Grande, RS, Brazil

¹⁶ Laboratório de Métodos de Estudos Subaquáticos e Cefalópodes, Departamento de Ecologia e Zoologia, Universidade Federal de Santa Catarina, Florianópolis, SC, Brazil

¹⁷ Departamento de Biologia Animal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, SP, Brazil

¹⁸ Laboratório de Ecologia de Manguezais (Ecomangue), Instituto Federal de Educação, Ciência e Tecnologia do Ceará, Acaraú, CE, Brazil

- ¹⁹ Laboratório de Moluscos, Centro de Pesquisa em Aquicultura e Pesca Sustentável do Semiárido, Departamento de Biociências, Universidade Federal Rural do Semiárido, Mossoró, RN, Brazil
- ²⁰ Departamento de Ciências Biológicas e Ambientais, Universidade Estadual Paulista Júlio de Mesquita Filho, Campus do Litoral Paulista, São Vicente, SP, Brazil
- ²¹ Departamento de Oceanografia Biológica, Instituto Oceanográfico, Universidade de São Paulo, São Paulo, Brazil
- ²² Grupo de Investigación y Desarrollo Tecnológico en Acuicultura y Pesca, Facultad Regional Chubut, Universidad Tecnológica Nacional, Puerto Madryn, Chubut, Argentina
- ²³ Setor de Malacologia, Departamento de Invertebrados, Museu Nacional, Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil
- ²⁴ Centro Universitario Regional del Este (CURE), Sede Rocha, Universidad de la República, Rocha, Uruguay
- ²⁵ Laboratório de Malacologia, Departamento de Zoologia, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brazil
- ²⁶ Centro de Ciências Computacionais, Universidade Federal do Rio Grande, Rio Grande, RS, Brazil
- ²⁷ Laboratório de Parasitologia—Malacologia, Instituto Butantan, São Paulo, SP, Brazil